

AN INDEX-CATALOGUE OF AFRICAN
PHANEROPTERINAE
(ORTHOPTERA : TETTIGONIIDAE)

BY

D. R. RAGGE, *cf.*
British Museum (Natural History)



Pp. 73-108

BULLETIN OF
THE BRITISH MUSEUM (NATURAL HISTORY)
ENTOMOLOGY Vol. 22 No. 3
LONDON : 1968

THE BULLETIN OF THE BRITISH MUSEUM
(NATURAL HISTORY), instituted in 1949, is
issued in five series corresponding to the Departments
of the Museum, and an Historical series.

Parts will appear at irregular intervals as they become ready. Volumes will contain about three or four hundred pages, and will not necessarily be completed within one calendar year.

In 1965 a separate supplementary series of longer papers was instituted, numbered serially for each Department.

This paper is Vol. 22, No. 3 of the Entomological series. The abbreviated titles of periodicals cited follow those of the World List of Scientific Periodicals.

World List abbreviation:
Bull. Br. Mus. nat. Hist. (Ent.).

© Trustees of the British Museum (Natural History) 1968

TRUSTEES OF
THE BRITISH MUSEUM (NATURAL HISTORY)

Issued 28 May, 1968

Price Sixteen Shillings

AN INDEX-CATALOGUE OF AFRICAN PHANEROPTERINAE (ORTHOPTERA : TETTIGONIIDAE)

By D. R. RAGGE

CONTENTS

| | <i>Page</i> |
|--|-------------|
| SYNOPSIS | 75 |
| INTRODUCTION | 75 |
| TREATMENT | 76 |
| THE STATUS OF THE PHANEROPTERINAE | 78 |
| CATALOGUE OF AFRICAN PHANEROPTERINAE | 79 |
| REFERENCES | 98 |
| INDEX | 103 |

SYNOPSIS

A catalogue is given of all the genera and species of Phaneropterinae known at present to occur in Africa and its off-shore islands (but not those of the Mediterranean Sea or the Malagasy Sub-region), and in the Arabian peninsula. The species are arranged alphabetically under the genera to which they have been assigned in the most recent literature, and the arrangement of the genera is also alphabetical. Generic and specific synonyms, junior homonyms and deliberate variations in spelling are listed chronologically under the valid name. The type-species is cited for all generic names, and full information on the type-material is given for all specific names. The catalogue includes 115 genus-group and 358 species-group names; of these, eighty-three generic, 295 specific and seven subspecific names are currently regarded as valid. The status of the Phaneropterinae is discussed.

INTRODUCTION

THIS catalogue has been compiled as the first stage of a comprehensive revision of the African Phaneropterinae. It includes all the species of Phaneropterinae known at present to occur in Africa, listed alphabetically under the genera to which they have been assigned in the most recent literature. As a deliberate policy, no new combinations (with the exception of an unavoidable one mentioned below) or synonyms are established in this paper; any necessary nomenclatorial changes will appear in the systematic revision to be published later. With the exception of groups covered by recent revisions (Ragge, 1955, 1956a, 1956b, 1957, 1960a, 1960b, 1961a, 1961b, 1962a, 1962b, 1964) the nomenclature will thus be subject to some alteration during the next few years. However, the standard of taxonomic work on the Tettigoniidae during most of the past century has been remarkably high and there are very few nomenclatorial problems.

The geographical region covered by this catalogue includes the whole of the African mainland and off-shore islands (e.g. Socotra, Zanzibar and the islands of the Gulf of Guinea) and the Arabian peninsula, but excludes the Malagasy Sub-region; there is

such a high degree of endemism in the Phaneropterinae of this sub-region that it is more appropriate to treat them separately. The Mediterranean islands are not included, and the inclusion of North Africa adds only thirteen species, all belonging to the genus *Odontura* Rambur.

TREATMENT

The eventual systematic arrangement of the African genera of Phaneropterinae will naturally depend on the comprehensive revision to be undertaken during the next few years, and it has therefore been thought better at this stage to list the genera alphabetically. Generic synonyms, junior homonyms and deliberate variations in spelling are listed chronologically under the valid generic name. The entries for each genus and generic synonym include a reference to the original description and the citation of the type-species and its mode of fixation; where the type-species was fixed by subsequent designation, a bibliographical reference is given to the fixation. The type-species is cited in its original combination; where it is now a junior synonym the present valid name is given in brackets. Where a type-species does not occur in Africa, the country of the type-locality is given in brackets. All generic synonyms are cited, including those based on non-African type-species; the type-species are cited in the same way as those of valid generic names.

The African species of each genus are listed alphabetically, together with references to their original descriptions and a statement of the nature and data of the type-material and the institution in which the type-material is housed. The absence of any information on the number and sex of the specimens in the type-series indicates that this information was not given with the original description and has not been determined subsequently. Where a species is based on a type-series from more than one locality, the citation of one depository after the last locality indicates that all the type-material is believed to be in that depository. It should be noted that the stated numbers of syntypes are not always known for certain, but always with at least a high degree of probability. A number of species described by Sjöstedt (1909) from Kilimanjaro and Meru were based on long series of syntypes, a proportion of which were sent out from Stockholm, where Sjöstedt worked, to other museums; as the exact extent of this distribution of syntypes is unknown, the only depository cited in these cases is the Naturhistoriska Riksmuseum, Stockholm.

Where a species has been transferred from its original genus, the name of the latter is given in brackets after the reference to the original description; a change in status is noted in a similar way. A question-mark before the original genus indicates that the original generic assignment was doubtful; one before the specific name itself indicates that the present generic assignment is doubtful. Where a species is currently regarded as polytypic, an entry is given for each subspecies and there is no separate entry for the species as a whole.

Specific synonyms (including those based on non-African type-specimens) and deliberate variations in spelling are indented and listed chronologically under each species or subspecies; they are otherwise given the same treatment as valid specific names. A question-mark before a specific synonym indicates that the synonymy is doubtful.

Territories in which type-localities are situated are given their current political names. Where the country is unknown the locality data originally published is quoted between inverted commas. Provinces are not cited unless given with the published locality data except in the case of South African specimens, for which the provinces (if known) are always cited.

Where a lectotype has been designated a bibliographical reference is given to the designation. No lectotypes are designated in this catalogue.

The museums and other institutions in which the type-material is housed are abbreviated in this catalogue as follows.

| | |
|-------|---|
| ANS | Academy of Natural Sciences of Philadelphia. |
| BMNH | British Museum (Natural History). |
| DEI | Deutsches Entomologisches Institut, Eberswalde (East Germany). |
| EIETH | Entomologisches Institut of the Eidgenössische Technische Hochschule, Zürich. |
| IEE | Instituto Español de Entomología, Madrid. |
| ILEA | Instituto e Laboratorio di Entomologia Agraria, Portici. |
| IMZU | Istituto e Museo di Zoologia of the Università di Torino, Turin. |
| IRSNB | Institut Royal des Sciences Naturelles de Belgique, Brussels. |
| IZPAN | Instytut Zoologiczny of the Polska Akademia Nauk, Warsaw. |
| IZU | Istituto di Zoologia of the Università di Pavia. |
| MCSN | Museo Civico di Storia Naturale, Genoa. |
| MHN | Muséum d'Histoire Naturelle, Geneva. |
| MLZA | Museu e Laboratório Zoológico e Antropológico, Lisbon. |
| MNHN | Muséum National d'Histoire Naturelle, Paris. |
| MRAC | Musée Royal de l'Afrique Centrale, Tervuren. |
| NM | Naturhistorisches Museum, Vienna. |
| NR | Naturhistoriska Riksmuseum, Stockholm. |
| SAM | South African Museum, Cape Town. |
| SMN | Staatliches Museum für Naturkunde, Stuttgart. |
| TM | Természettudományi Múzeum, Budapest. |
| UM | University Museum, Oxford. |
| UZI | Universitetets Zoologiska Institution, Lund. |
| ZM | Zoologisches Museum, Hamburg. |
| ZMHU | Zoologisches Museum of the Humboldt-Universität, Berlin. |

In the course of compiling this catalogue I have come across one generic and two specific names published before Kirby's catalogue of 1906, but not included in it. The generic name is *Cameronia*, which first appeared in the combination *Symmetropleura (Cameronia) africana* Brunner in a paper by Karsch (1889 : 450). The name was ignored by Brunner (1891) in the supplement to his earlier monograph (Brunner, 1878) but was quoted, in the same combination, by Bolívar (1906 : 349). As far as I know these are the only two occasions on which this name has appeared in print. *Cameronia* is clearly a genus-group name and is available under Article 16 (a) (v) of the *International Code of Zoological Nomenclature*, with *Symmetropleura africana* Brunner as its type-species, by monotypy. It seems highly improbable that Karsch intended the name to be regarded as a subgenus of *Symmetropleura* Brunner; there is no other instance of either him or his contemporary Brunner using a subgeneric name, and on the following page of his 1889 paper (*op. cit.*, p. 451) he describes another

species of *Symmetropleura* without giving it a subgeneric assignment. It seems more likely that Karsch was considering *Cameronia* as an alternative generic name for *Symmetropleura africana* Brunner, and so cited it in brackets. No subgenera are recognized in this catalogue and I have merely listed *Cameronia* in the synonymy of *Symmetropleura*. If, as seems probable, revisionary work shows that the African species at present assigned to the otherwise New World genus *Symmetropleura* should be separated at the generic level, the name *Cameronia* is available for any resulting genus that contains *Symmetropleura africana* Brunner.

The two specific names missed by Kirby (1906) are *Cestromoecha crassipes* Karsch (described in a footnote—Karsch, 1890 : 364) and *Rhegmatopoda brunneri* Kirby. The latter name, proposed as a nomen novum (Kirby, 1900 : 215), was based on Brunner's (1878 : 89) description of a male specimen from Grahamstown in the Naturhistorisches Museum, Vienna, then regarded by Brunner as *Horatospaga leptocerca* Stål, and Kirby's examination of a male specimen from Barberton in the British Museum (Natural History). Although these specimens were not described by Kirby, the name *R. brunneri* is available under Article 16 (a) (i) of the *International Code of Zoological Nomenclature*. The name *Rhegmatopoda* Brunner has since become a junior synonym of *Conchotopoda* Karsch (Ragge, 1960a : 325) and *brunneri* is thus listed in this catalogue under *Conchotopoda* Karsch, forming the new combination *Conchotopoda brunneri* (Kirby) **comb. n.**

THE STATUS OF THE PHANEROPTERINAE

The inter-relationships and systematic arrangement of the African genera of Phaneropterinae will be considered in the course of forthcoming revisionary work, but it is appropriate at this stage to consider briefly the status of the group as a whole.

The Phaneropterinae have been regarded as a subfamily of Tettigoniidae (also called at various times, and often in a broader sense, Locustariae, Locustina, Locustidae, Locustodea and Phasgonuridae) for well over a century: they were given this status, together with *Mecopoda* and *Phyllophora* (now the type-genera of two other subfamilies), by Burmeister (1838 : 684). In 1878 Brunner, in his classic *Monographie der Phaneropteriden*, established the group in its present sense and laid the foundation for all subsequent systematic work on the Phaneropterinae. The subfamily suffix *-inae* was not in use at this time, but Brunner referred to the group as a "Subfamilie" and used for it the Latin category "tribus". Brunner reaffirmed his classification in his works of 1891 and 1893. The group-name appeared with the currently used subfamily suffix in Saussure & Pictet's work of 1897, and was again used in this form by Kirby (1906). This usage was adopted almost universally and still prevails today (Bei-Bienko, 1954, 1964; Blatchley, 1920; Brues, Melander & Carpenter, 1954; Chopard, 1922, 1943, 1947; Essig, 1947; Grant, 1964; Imms, 1957; Zeuner, 1936a).

During this period the general classification of the Ensifera came under the strong stabilizing influence of two major works on their morphology and phylogeny, one by Ander and the other by Zeuner, published almost simultaneously in 1939. The thoroughness and comprehensiveness of these studies clearly makes them the best available basis for considering the inter-relationships of the sub-groups of Ensifera,

and it is significant that the classifications adopted in the two works are almost identical. The Phaneropterinae were again regarded as a subfamily of the Tettigoniidae.

This long period of stability in the status of the Phaneropterinae has been affected during the past twenty years by a movement among European orthopterists to raise the status of nearly all the groups of Saltatoria, i.e. to elevate most of the families to superfamilies or even suborders, and most of the subfamilies to families. This movement does not stem from new morphological or phylogenetic discoveries but simply from the view that these groups are at present ranked at too low a level, or, in one case (Dirsh, 1961 : 358), from the belief that family and subfamily rank can be objectively defined. The effect of this action on the Phaneropterinae is to raise them to family rank, the Tettigoniidae becoming a superfamily more restricted in scope than the Tettigonioidea of Ander (*op. cit.*), which is left without a name.

Although there are a few earlier instances of the Phaneropterinae being accorded family status (e.g. Burr, 1910; Zacher, 1917), the first major recent work in which this was done was that of Chopard (1949), and the practice has since been adopted by Beier (1955) and in the faunal works of Harz (1957, 1960). Although it seems likely that some other modern orthopterists whose specialities lie outside the Ensifera would favour family status for the Phaneropterinae, the group is still regarded as a subfamily in most recent taxonomic publications in both Europe and America (e.g. Bei-Bienko, 1964; Grant, 1964; Karabag, 1964; Ragge, 1964); indeed, some orthopterists (e.g. Uvarov, 1924 : 492–493; Zeuner, 1936a : 107, 1936b : 300–301, 1939 : 115) have suggested that a number of the other subfamilies of Tettigoniidae would be better regarded as tribes.

There is of course a trend during the earlier stages of the taxonomic study of all groups towards the raising in rank of each level of the hierarchy, and it may seem at first sight that there is still a need for such a trend in the classification of the Ensifera. However, in my view the classification firmly established by Ander (*op. cit.*) and since widely adopted by orthopterists offers an acceptable compromise in the rank accorded to each group, and there is no reason why it should not be used, with minor modifications, as a framework for taxonomic work on this suborder for many years to come. In the case of the Phaneropterinae there is room for only one category between the group as a whole and its genera, one that is well provided for by tribes; if the group is regarded as a family, either the subfamily or tribe category will be left unemployed. This is not the place for further discussion of the problems of the classification of the Ensifera; it suffices for me to lend my support to the view still expressed by the majority of orthopterists—that the Phaneropterinae are best regarded as a subfamily.

CATALOGUE OF AFRICAN PHANEROPTERINAE

PHANEROPTERINAE Burmeister

Phaneropteridae Burmeister, 1838 : 684.

Camptoxiphiae Serville, [1838] : 399. (Not based on the name of a contained genus and therefore unavailable under Article 11 (e) of the *International Code of Zoological Nomenclature*.)

Phaneropterinae; Saussure & Pictet, 1897 : 310.

ANISOTOCHRA Karsch

Anisotochra Karsch, 1889 : 430. Type-species: *Anisotochra gracilipes* Karsch, by monotypy.
gracilipes Karsch, 1889 : 431. Holotype ♀, CONGO (KINSHASA): Kuako to Kimpoko (ZMHU, Berlin).

ARANTIA Stål

Arantia Stål, 1874 : 25. Type-species: *Holochlora fatidica* Stål, by monotypy.

accrana Karsch, 1889 : 436. Holotype ♀, GHANA: Accra (ZMHU, Berlin).
angustipennis Chopard, 1954 : 36. Holotype ♂, GUINEA: Nimba (MNHN, Paris).
brevipes Chopard, 1954 : 37. 1 ♂, 1 ♀ syntypes, GUINEA: Nimba (MNHN, Paris).
excelsior Karsch, 1889 : 435. Holotype ♂, SIERRA LEONE (IZPAN, Warsaw).
fasciata Walker, 1869 : 378 (*Phylloptera*). Holotype ♀, GAMBIA (BMNH, London).
atrolineata Brunner, 1891 : 69. Holotype ♀, "SLAVE COAST" (EIETH, Zürich).
fatidica Stål, 1873 : 43 (*Holochlora*). Holotype ♂, SIERRA LEONE (NR, Stockholm).
gabunensis Brunner, 1891 : 64. Holotype ♀, GABON (NM, Vienna).
gestri Griffini, 1906 : 384. Holotype ♂, CONGO (BRAZZAVILLE): N'jole (MCSN, Genoa).
hydatinoptera Karsch, 1889 : 436. 1 ♂, 1 ♀ syntypes, NIGERIA (IZPAN, Warsaw).
latifolia Karsch, 1890c : 356. 1 ♂, 1 ♀ syntypes, CAMEROUN : Barombi (ZMHU, Berlin).
leptocnemis Karsch, 1890c : 358. Holotype ♀, CAMEROUN : Barombi (ZMHU, Berlin).
mammisignum Karsch, 1896 : 332. Holotype ♀, CAMEROUN: Lolodorf (ZMHU, Berlin).
manca Bolívar, 1906 : 332. Holotype ♂, CAMEROUN (IEE, Madrid).
marmorata Karsch, 1889 : 435. Holotype ♀, "CONGO" (IZPAN, Warsaw).
melanotus Sjöstedt, 1901 : 6. Holotype ♂, CAMEROUN: Bonge (NR, Stockholm).
orthocnemis Karsch, 1890c : 357. 1 ♂ syntype, CAMEROUN; 1 ♀ syntype, CAMEROUN: Barombi (ZMHU, Berlin).
ovalipennis Chopard, 1954 : 38. Holotype ♀, GUINEA: Nimba, N'Zo (MNHN, Paris).
rectifolia Brunner, 1878 : 137. Holotype ♂, FERNANDO POO (NM, Vienna).
regina Karsch, 1888 : 434. Holotype ♀, GABON (ZMHU, Berlin).
retinervis Karsch, 1888 : 437. 2 ♂ syntypes, CAMEROUN: Barombi (ZMHU, Berlin); 1 ♂ syntype, "Chinchoco" (ZMHU, Berlin); 1 ♂ syntype, SIERRA LEONE (IZPAN, Warsaw).
scurra Karsch, 1896 : 331. Holotype ♂, CAMEROUN: Lolodorf (ZMHU, Berlin).
simplicinervis Karsch, 1889 : 438. 1 ♂ syntype, "Chinchoco"; 1 ♀ syntype, CONGO (KINSHASA): Kuako to Kimpoko (ZMHU, Berlin).
spinulosa Brunner, 1878 : 137. 1 ♂, 2 ♀ syntypes, SOUTH AFRICA: Natal, Durban (NM, Vienna).
tigrina Bolívar, 1906 : 330. Holotype ♂, FERNANDO POO (IEE, Madrid).
ugandana Rehn, 1914 : 176. Holotype ♀, UGANDA (DEI, Eberswalde).

ATLASACRIS Rehn

Atlasacris Rehn, 1914 : 153. Type-species: *Atlasacris peculiaris* Rehn, by original designation.
peculiaris Rehn, 1914 : 154. Holotype ♂, "Ost-AFRIKA" (ZMHU, Berlin).

AZAMIA Bolívar

Azamia Bolívar, 1906 : 339. Type-species: *Azamia biplagiata* Bolívar, by monotypy.

biplagiata Bolívar, 1906 : 341. Holotype ♂, CAMEROUN (IEE, Madrid).
doriae Griffini, 1906 : 395 (*Vossia*). Holotype ♂, FERNANDO POO: Musola (MCSN, Genoa).

BONGEIA Sjöstedt

Bongeia Sjöstedt, 1901 : 9. Type-species: *Bongeia puncticollis* Sjöstedt, by monotypy.

brevicauda Ebner, 1943 : 265. Holotype ♀, FERNANDO POO (DEI, Eberswalde).
puncticollis Sjöstedt, 1901 : 9. Holotype ♀, CAMEROUN (NR, Stockholm).

BUEACOLA Sjöstedt

Bueacola Sjöstedt, 1912 : 7. Type-species: *Bueacola cornigera* Sjöstedt, by monotypy.

cornigera Sjöstedt, 1912 : 8. Holotype ♂, CAMEROUN: Buea (NR, Stockholm).

BUETTNERIA Karsch

Büttneria Karsch, 1889 : 444. Type-species: *Buettneria maculiceps* Karsch, by monotypy.

Buettneria Karsch, Brunner, 1891 : 105.

Stenacropteryx Karsch, 1896 : 339. Type-species: *Stenacropteryx eburneiguttata* Karsch
 (= *Buettneria maculiceps* Karsch), by monotypy.

maculiceps Karsch, 1889 : 445. Holotype ♀, CONGO (KINSHASA): Kuako to Kimpoko (ZMHU, Berlin).

eburneiguttata Karsch, 1896 : 339 (*Stenacropteryx*). Holotype ♂, CAMEROUN: Lolodorf (ZMHU, Berlin).

CATOPTROPTERYX Karsch

Catoptropteryx Karsch, 1890c : 361. Type-species: *Catoptropteryx guttatipes* Karsch, by monotypy (two other species doubtfully included).

africana Karsch, 1889 : 446 (*Caedicia*). Holotype ♀, NIGERIA: Benue (IZPAN, Warsaw).

apicalis Bolívar, 1893 : 177 (*Caedicia*). Holotype ♀, IVORY COAST: Assinie (IEE, Madrid).

capreola Karsch, 1896 : 332. Holotype ♂, CAMEROUN: Lolodorf (ZMHU, Berlin).

extensipes Karsch, 1896 : 336. Holotype ♂, CAMEROUN: Lolodorf (ZMHU, Berlin).

guttatipes Karsch, 1890c : 362. Holotype ♂, CAMEROUN: Barombi (ZMHU, Berlin).

immaculipennis Karsch, 1896 : 333. Holotype ♂, CAMEROUN: Lolodorf (ZMHU, Berlin).

latipennis Chopard, 1955 : 267. Holotype ♂, SOUTH AFRICA: Cape Province, Tzitzikama Forest, Stormsrivierpiek (UZI, Lund).

maculipennis Karsch, 1896 : 333. Holotype ♀, CAMEROUN: Lolodorf (ZMHU, Berlin).

neutralipennis Karsch, 1896 : 334. 1 ♂ syntype, TOGO: Misahöhe; 1 ♀ syntype, TOGO: Bismarckburg (ZMHU, Berlin).

nigrospinosa Brunner, 1891 : 97 (*Caedicia*). Holotype ♀, CAMEROUN (IZPAN, Warsaw).

punctulata Karsch, 1890b : 260 (*Caedicia*). Holotype ♀, CAMEROUN: Kribi (ZMHU, Berlin).

signatipennis Karsch, 1896 : 334. 1 ♂ syntype, CAMEROUN: Victoria; 1 ♀ syntype, CAMEROUN: Lolodorf (ZMHU, Berlin).

CESTROMOECHA Karsch

Cestromoecha Karsch, 1893 : 128. Type-species: *Poreuomena tenuipes* Karsch, by original designation.

crassipes Karsch, 1890c : 364 (*Poreuomena*). Holotype ♀, CAMEROUN (ZMHU, Berlin).

mundamensis Karsch, 1896 : 338. 1 ♂, 1 ♀ syntypes, CAMEROUN: Mundame (ZMHU, Berlin).

tenuipes Karsch, 1890c : 363 (*Poreuomena*). 3 ♀ syntypes, CAMEROUN: Barombi (ZMHU, Berlin).

CONCHOTOPODA Karsch

- Conchotopoda* Karsch, 1887a : 44. Type-species: *Conchotopoda belcki* Karsch, by monotypy.
Rhegmatopoda Brunner, 1891 : 44. Type-species: *Horatosphaga leptocerca* Stål, by monotypy.
- belcki* Karsch, 1887a : 45. Holotype ♂, SOUTH WEST AFRICA: Damaraland (ZMHU, Berlin).
brunneri Kirby, 1900 : 215 (*Rhegmatopoda*) (see p. 78). 1 ♂ syntype, SOUTH AFRICA: Transvaal, Barberton (BMNH, London); 1 ♂ syntype, SOUTH AFRICA: Cape Province, Grahams-town (NM, Vienna).
- crassicauda* Ragge, 1960a : 328. Holotype ♂, SOUTH AFRICA: Transvaal, Pietersburg (SAM, Cape Town).
- grallatoria* Stål, 1856 : 170 (?*Phaneroptera*). Holotype ♂, SOUTH AFRICA: "Caffraria" (NR, Stockholm).
- leptocerca* Stål, 1876 : 59 (*Horatosphaga*). Holotype ♂, SOUTH WEST AFRICA: Damara (NR, Stockholm).
- parva* Ragge, 1960a : 330. Holotype ♂, SOUTH AFRICA: Orange Free State, Witzieshoek (BMNH, London).

CORYCOMIMA Karsch

- Corycomima* Karsch, 1889 : 457 (proposed conditionally, but available from this date under Article 17 (8) of the International Code of Zoological Nomenclature). Type-species: *Plangia camerata* Karsch (= *Corycomima flavescentis* (Walker)), by monotypy.
- Corycomima* Karsch, 1896 : 343 (proposed unconditionally).

- flavescentis* Walker, 1869 : 386 (*Orophus*). Holotype ♀, GAMBIA (BMNH, London).
camerata Karsch, 1889 : 457 (*Plangia*). Holotype ♀, CAMEROUN: Barombi (ZMHU, Berlin).

CORYMETA Brunner

- Corymeta* Brunner, 1878 : 126. Type-species: *Phaneroptera amplexens* Schaum, by monotypy.
- amplexens* Schaum, 1853 : 777 (*Phaneroptera*). Holotype ♂, MOZAMBIQUE (ZMHU, Berlin).

DANNFELTIA Sjöstedt

- Dannfeltia* Sjöstedt, 1901 : 19. Type-species: *Dannfeltia nana* Sjöstedt, by monotypy.
- nana* Sjöstedt, 1901 : 19. Holotype ♂, "KONGO" (NR, Stockholm).

DAPANERA Karsch

- Dapanera* Karsch, 1889 : 441. Type-species: *Dapanera genuteres* Karsch, by monotypy.
- eidmanni* Ebner, 1943 : 263. Holotype ♂, FERNANDO Poo (DEI, Eberswalde).
- genuteres* Karsch, 1889 : 441. 2 ♂ syntypes, GHANA: Accra (ZMHU, Berlin).
- irregularis* Karsch, 1890b : 258. 1 ♂, 1 ♀ syntypes, CAMEROUN: Kribi (ZMHU, Berlin).

DEBRONA Walker

- Debrona* Walker, 1870 : 480. Type-species: *Debrona cervina* Walker, by monotypy.
- Otiaphysa* Karsch, 1889 : 453. Type-species: *Otiaphysa habitata* Karsch (= *Debrona cervina* Walker), by monotypy.
- angustipennis* Burr, 1900 : 43 (*Otiaphysa*). Holotype ♀, SOMALI REPUBLIC: Whardi Datal (UM, Oxford).
- cervina* Walker, 1870 : 481. Holotype ♀, SOUTH AFRICA (BMNH, London).
- habitata* Karsch, 1889 : 454 (*Otiaphysa*). Holotype ♀, TANZANIA: Usambara (ZMHU, Berlin).

DIOGENA Brunner

Diogena Brunner, 1878 : 224. Type-species: *Phaneroptera fausta* Burmeister, by monotypy.

denticulata Chopard, in Chopard & Kevan, 1954 : 328. Holotype ♀, KENYA: Northern Frontier Distr., El Wak (BMNH, London).

fausta Burmeister, 1838 : 689 (*Phaneroptera*). Holotype ♂, SUDAN: Nubia (ZMHU, Berlin).

DIONCOMENA Brunner

Dioncomena Brunner, 1878 : 208. Type-species: *Dioncomena ornata* Brunner, by monotypy.

ornata Brunner, 1878 : 208. 2 ♂, 1 ♀ syntypes, "ZANZIBAR" (NM, Vienna); 1 ♂ syntype, "ZANZIBAR" (IEE, Madrid).

superba Karsch, 1889 : 449. Unknown number of syntypes of both sexes, TANZANIA: Bondei and Usambara (ZMHU, Berlin).

DITHELA Karsch

Dithela Karsch, 1890c : 354. Type-species: *Dithela rectiloba* Karsch, by monotypy.

acuticercus Sjöstedt, 1912 : 5. 2 ♂, 1 ♀ syntypes, CAMEROUN: Buea (NM, Stockholm).

rectiloba Karsch, 1890c : 355. 1 ♂, 2 ♀ syntypes, CAMEROUN: Barombi (ZMHU, Berlin).

DREPANO PHYLLUM Karsch

Drepanophyllum Karsch, 1890c : 358. Type-species: *Drepanophyllum marmoratum* Karsch, by monotypy.

Karschia Brunner, 1891 : 41. Type-species: *Karschia corrosa* Brunner (= *Drepanophyllum marmoratum* Karsch), by monotypy.

corrosifolium Karsch, 1896 : 326. Unknown number of syntypes of both sexes, CAMEROUN: Yaunde and Lolodorf (ZMHU, Berlin).

furcatum Ragge, 1962a : 307. Holotype ♂, UGANDA: Entebbe (BMNH, London).

marmoratum Karsch, 1890c : 360. Holotype ♀, CAMEROUN: Barombi (ZMHU, Berlin).

corrosa Brunner, 1891 : 41 (*Karschia*). Holotype ♀, CAMEROUN (IZPAN, Warsaw).

DUCETIA Stål

Ducetia Stål, 1874 : 11. Type-species: *Locusta japonica* Thunberg, by monotypy (type-locality: Japan).

Paura Karsch, 1889 : 439. Type-species: *Paura biramosa* Karsch, by subsequent designation (Kirby, 1906 : 407).

Pseudisotima Schulthess, 1898 : 199. Type-species: *Pseudisotima punctata* Schulthess, by monotypy.

Kuwayamaea Matsumura & Shiraki, 1908 : 7. Type-species: *Kuwayamaea sapporensis* Matsumura & Shiraki (= *Ducetia chinensis* (Brunner)), by original designation (type-locality: Japan).

Telaea Bolívar, 1922 : 201. Type-species: *Telaea quadripunctata* Bolívar (= *Ducetia punctipennis* (Gerstaecker)), by monotypy.

biramosa Karsch, 1889 : 439 (*Paura*). Holotype ♀, TANZANIA: Usambara (ZMHU, Berlin).

chelocerca Ragge, 1961a : 197. Holotype ♂, SOUTH AFRICA: Transvaal, junction of Crocodile and Marico Rivers (SAM, Cape Town).

- costata* Ragge, 1961a : 196. Holotype ♂, CONGO (KINSHASA): Nyangwe (MRAC, Tervuren).
- crosskeyi* Ragge, 1961a : 192. Holotype ♂, NIGERIA: Niger Province, Diko (BMNH, London).
- fuscopunctata* Chopard, 1954 : 35. Holotype ♂, GUINEA: Nimba, Yalanzou (MNHN, Paris).
- loosi* Griffini, 1908b : 204. Holotype ♀, CONGO (KINSHASA): Popocabacca (IRSNB, Brussels).
- macrocerca* Ragge, 1961a : 195. Holotype ♂, CONGO (KINSHASA): Katanga, Kiambi (MRAC, Tervuren).
- parva* Ragge, 1961a : 205. Holotype ♂, SOMALI REPUBLIC: Haud (BMNH, London).
- punctata* Schulthess, 1898 : 199 (*Pseudisotima*). Holotype ♀, ETHIOPIA: Girma (MCSN, Genoa).
- punctipennis* Gerstaecker, 1869 : 215 (*Phaneroptera*). Holotype ♂, KENYA: Mt. Ndara (ZMHU, Berlin).
- reticulosa* Karsch, 1889 : 440 (*Paura*). Holotype ♀, KENYA: Mombasa (ZMHU, Berlin).
- quadripunctata* Bolívar, 1922 : 201 (*Telaea*). Holotype ♂, KENYA: Simba (lost).
- ramulosa* Ragge, 1961a : 198. Holotype ♂, ZAMBIA: Luano Valley, R. Mulungushi (BMNH, London).
- sagitta* Ragge, 1961a : 199. Holotype ♂, ANGOLA: Lepi (BMNH, London).
- vitrialia* Ragge, 1961a : 204. Holotype ♂, SOMALI REPUBLIC: Haud (BMNH, London).

ENOCHLETICA Karsch

- Enochletica* Karsch, 1896 : 336. Type-species: *Enochletica ostentatrix* Karsch, by monotypy.
- ostentatrix* Karsch, 1896 : 337. Unknown number of syntypes of both sexes, CAMEROON: Lolodorf (ZMHU, Berlin).

EPIPHLEBUS Karsch

- Epiphlebus* Karsch, 1896 : 325. Type-species: *Epiphlebus crypterius* Karsch, by monotypy.
- crypterius* Karsch, 1896 : 325. Holotype ♀, SOMALI REPUBLIC (type-depository unknown).
- ruspolii* Schulthess, 1898 : 197. Holotype ♀, SOMALI REPUBLIC: Ganana (MCSN, Genoa).

EULIOPTERA Ragge

- Eulioptera* Ragge, 1956a : 266. Type-species: *Phaneroptera reticulata* Brunner, by original designation.
- breviala* Ragge, 1956a : 277. Holotype ♂, CONGO (KINSHASA): Kivu, Kibumba (MRAC, Tervuren).
- longicerca* Ragge, 1956a : 273. Holotype ♂, CONGO (KINSHASA): Kinshasa (MRAC, Tervuren).
- reticulata leptomorpha* Ragge, 1956a : 271. Holotype ♂, ZAMBIA: Mporokoso distr., Mweru Wa Ntipa (BMNH, London).
- reticulata reticulata* Brunner, 1878 : 213 (*Phaneroptera reticulata*). Holotype ♀, SOUTH AFRICA: Cape Province, Grahamstown (NM, Vienna).
- spinulosa* Ragge, 1956a : 274. Holotype ♂, CONGO (KINSHASA): Kasenyi (BMNH, London).

EURYCORYPHA Stål

- Eurycorypha* Stål, 1873 : 40. Type-species: *Phylloptera cereris* Stål, by original designation.
- Myrmecophana* Brunner, 1883 : 248. Type-species: *Myrmecophana fallax* Brunner, by monotypy.

- adicra* Karsch, 1892b : 75. Holotype ♀, CAMEROON: Buea (ZMHU, Berlin).
- aequatorialis* Krauss, 1890 : 663. Holotype ♀, ROLAS I. (Gulf of Guinea) (SMN, Stuttgart).

- arabica arabica** Uvarov, 1936 : 534. Holotype ♂, SAUDI ARABIA: S. Hedjaz, Taif (BMNH, London).
- arabica media** Uvarov, 1936 : 535. 3 ♂ syntypes, SAUDI ARABIA: El Kubar (BMNH, London).
- arabica reducta** Uvarov, 1936 : 536. Holotype ♂, SOUTH ARABIA: Hadramaut, Wadi Humum, nr. Makalla (ZM, Hamburg).
- brevicollis** Stål, 1876 : 61. Holotype ♀, SOUTH WEST AFRICA: Ovambo (NR, Stockholm).
- canaliculata** Karsch, 1890b : 261. Holotype ♂, CAMEROUN: Kribi (ZMHU, Berlin).
- cereris** Stål, 1856 : 170 (*Phylloptera*). Holotype ♂, SOUTH AFRICA: Natal, Durban (NR, Stockholm).
- compressa** Walker, 1869 : 386 (*Orophus*). Holotype ♂, SOUTH AFRICA (BMNH, London).
- cuspidata** Krauss, 1901 : 292. 1 ♂ syntype, SOUTH WEST AFRICA: Okahandja; 1 ♂ syntype, SOUTH WEST AFRICA: Windhoek (SMN, Stuttgart).
- darlingi** Uvarov, 1936 : 536. Holotype ♂, SOUTH YEMEN: Makhnun (BMNH, London).
- diminuta** Chopard, 1938 : 121. Holotype ♂, KENYA: Turkana Desert, Lodwar (MNHN, Paris).
- fallax** Brunner, 1883 : 248 (*Myrmecophana*). Nymphal holotype, sex unknown, SUDAN: Ambucarra (IRSNB, Brussels).
- montana** Sjöstedt, 1901. Holotype ♀, CAMEROUN: Mapanja (NR, Stockholm).
- mutica** Karsch, 1892 : 327. Holotype ♀, CAMEROUN: Barombi (ZMHU, Berlin).
- ornatipes** Karsch, 1890b : 260. Holotype ♂, CAMEROUN: Kribi (ZMHU, Berlin).
- prosperinae** Brunner, 1878 : 274. Holotype ♀, SOUTH AFRICA: Natal, Durban (NM, Vienna).
- punctipennis** Chopard, 1938 : 121. Holotype ♀, KENYA: Mt. Elgon (MNHN, Paris).
- securifera** Brunner, 1878 : 275. Holotype ♂, SOUTH WEST AFRICA: Herrero-Land (NM, Vienna).
- simillima** Chopard, in Chopard & Kevan, 1954 : 330. Holotype ♂, KENYA: N. Turkana, Lokitaung (BMNH, London).
- spinulosa** Karsch, 1889 : 455. Holotype ♀, CONGO (KINSHASA): Kuako to Kimpoko (ZMHU, Berlin).
- strangulata** Walker, 1869 : 385 (*Orophus*). 1 ♀ syntype, SIERRA LEONE; 1 ♀ syntype, "WEST AFRICA" (BMNH, London).
- stylata** Stål, 1873 : 43. Holotype ♂, SIERRA LEONE (NR, Stockholm).
- sudanensis** Giglio-Tos, 1907 : 18. Holotype ♂, SUDAN: Port Sudan (IMZU, Turin).
- varia** Brunner, 1891 : 136. 1 ♂, 1 ♀ syntypes, TANZANIA: Kilimanjaro (NM, Vienna).
- velicauda** Karsch, 1893 : 131. 1 ♂, 1 ♀, 1 nymphal syntypes, TOGO: Bismarckburg (ZMHU, Berlin).
- zebrata** Bruner, 1920 : 135. Holotype ♀, CAMEROUN: Lolodorf (ANS, Philadelphia).

GABONELLA Uvarov

- Gabonia** Bolívar, 1906 : 327. Type-species: *Gabonia cothurnata* Bolívar, by monotypy.
(Homonym of *Gabonia* Jacoby, 1893 : 101.)
- Gabonella** Uvarov, 1940a : 174 (proposed as replacement name for *Gabonia* Bolívar).

cothurnata Bolívar, 1906 : 328 (*Gabonia*). 2 ♂, 3 ♀ syntypes, CAMEROUN (IEE, Madrid).

GELATOPOIA Brunner

- Gelatopoia** Brunner, 1891 : 111. Type-species: *Gelatopoia bicolor* Brunner, by monotypy.
- Gelatopoea** Brunner, Bolívar, 1906 : 349.
- Gelatopoea** Kirby, 1906 : 439. (Unjustified emendation.)

bicolor Brunner, 1891 : 112. Holotype ♂, SIERRA LEONE (IZPAN, Warsaw).

GOETIA Karsch

Goëtia Karsch, 1892a : 322. Type-species: *Goetia galbana* Karsch, by monotypy.
Goetia Karsch, Kirby, 1906 : 428.

dimidiata Bolívar, 1906 : 335. 1 ♂, 1 ♀ syntypes, CAMEROUN; 1 ♂, 1 ♀ syntypes, FERNANDO Poo; 1 ♂, 1 ♀ syntypes, no data (IEE, Madrid).

galbana Karsch, 1892a : 323. Holotype ♀, CAMEROUN: Barombi (ZMHU, Berlin).

GONATOXIA Karsch

Gonatoxia Karsch, 1889 : 441. Type-species: *Gonatoxia immaculata* Karsch, by subsequent designation (Kirby, 1906 : 424).

immaculata Karsch, 1889 : 442. Holotype ♀, TANZANIA: Usambara (ZMHU, Berlin).
maculata Karsch, 1889 : 442. Holotype ♂, SOMALI REPUBLIC (IZPAN, Warsaw).

GRAVENREUTHIA Karsch

Gravenreuthia Karsch, 1892b : 73. Type-species: *Gravenreuthia saturata* Karsch, by monotypy.

saturata Karsch, 1892b : 74. 1 ♂, 2 ♀ syntypes, CAMEROUN : Buea (ZMHU, Berlin).

HARPOSCEPA Karsch

Harposcepa Karsch, 1896 : 328. Type-species: *Harposcepa lobulipennis* Karsch, by monotypy. 1896.

karschiana Schulthess, 1898 : 202. Holotype ♀, SOMALI REPUBLIC (or ETHIOPIA?): Gubbet (IZU, Pavia).

lobulipennis Karsch, 1896 : 329. Holotype ♀, SOMALI REPUBLIC (type-depository unknown).

HIMERTULA Uvarov

Himerta Brunner, 1878 : 118. Type-species: *Himerta marginata* Brunner, by monotypy (type-locality: India). (Homonym of *Himerta* Foerster, 1868 : 200.)

Himertula Uvarov, 1940a : 174 (proposed as replacement name for *Himerta* Brunner).

feana Griffini, 1906 : 380 (*Himerta*). Holotype ♂, CONGO (BRAZZAVILLE): Nkogo (MCSN, Genoa).

HORATOSPHAGA Schaum

Horatosphaga Schaum, 1853 : 778. Type-species: *Horatosphaga serrifera* Schaum, by monotypy.

Plegmatoptera Karsch, 1889 : 429. Type-species: *Plegmatoptera reticulata* Karsch, by monotypy. (Homonym of *Plegmatoptera* Spinola, 1839 : 283.)

Pantolepta Karsch, 1889 : 429. Type-species: *Pantolepta heteromorpha* Karsch, by monotypy.

Pachypyga Karsch, 1893 : 124. Type-species: *Pachypyga inclusa* Karsch, by monotypy. (Homonym of *Pachypyga* Steinheil, 1873 : 564.)

Euthypphlebia Schulthess, 1898 : 204. Type-species: *Euthypphlebia parallela* Schulthess (= *Horatosphaga heteromorpha* (Karsch)), by monotypy.

Thaumatoxenia Kirby, 1909 : 65. Type-species: *Thaumatoxenia leggei* Kirby, by monotypy.

Keniacola Sjöstedt, 1912 : 13. Type-species: *Keniacola gracilis* Sjöstedt, by monotypy.

Schulthessinia Sjöstedt, 1912 : 15. Type-species: *Conchotopoda ruspolii* Schulthess, by original designation.

Pachypygiana Strand, 1921 : 106 (proposed as replacement name for *Pachypyga* Karsch).

Eupantolepta Bolívar, 1922 : 200. Type-species: *Eupantolepta regularis* Bolívar, by monotypy.

Plegmatia Uvarov, 1940b : 116 (proposed as replacement name for *Plegmatoptera* Karsch).

concava Ragge, 1960 : 308. Holotype ♂, UGANDA: Gulu (BMNH, London).

crosskeyi Ragge, 1960 : 318. Holotype ♂, NIGERIA: Niger Province, Diko (BMNH, London).

diminuta Chopard, in Chopard & Kevan, 1954 : 319 (*Peronura*). Holotype ♂, KENYA: Moyale (BMNH, London).

elgonis Chopard, 1938 : 119 (*Peronura*). Holotype ♂, KENYA: Mt. Elgon (MNHN, Paris).

elongata Rehn, 1914 : 163 (*Conchotopoda*). Holotype ♂, CONGO (KINSHASA): Usumbura (ZMHU, Berlin).

? *kandti* Rehn, 1914 : 167 (? *Horatosphaga*). Holotype ♀, CONGO (KINSHASA): Lake Kivu (ZMHU, Berlin).

gracilis Sjöstedt, 1912 : 14 (*Keniacola*). Holotype ♂, KENYA: Mt. Kenya (NR, Stockholm)

heteromorpha Karsch, 1889 : 430 (*Pantolepta*). Lectotype ♂, KENYA: Mombasa (ZMHU, Berlin); designated by Ragge (1960 : 298).

hoehneli Brunner, 1891 : 44 (*Plegmatoptera*). Holotype ♀, TANZANIA: Kilimanjaro (NM, Vienna).

parallela Schulthess, 1898 : 205 (*Euthyphlebia*). Holotype ♂, ETHIOPIA: Dolo (MCSN, Genoa).

kilimandjarica Sjöstedt, 1909 : 132 (*Pantolepta*). Lectotype ♂, TANZANIA: Mt. Meru (NR, Stockholm); designated by Ragge (1960 : 298).

gracilis Bolívar, 1922 : 198 (*Plegmatoptera*). Holotype ♂, KENYA: Loroghi Mtns. (MNHN, Paris).

inclusa Karsch, 1893 : 125 (*Pachypyga*). 1 ♂, 5 ♀ syntypes, TOGO: Bismarckburg (ZMHU, Berlin).

leggei Kirby, 1909 : 65 (*Thaumatoxenia*). Lectotype ♂, UGANDA: E. Ruwenzori, Mubuku Valley (BMNH, London); designated by Ragge (1960 : 292).

robusta Rehn, 1914 : 156. Holotype ♀, CONGO (KINSHASA): 90 km. W. of Albert-Nyanza (ZMHU, Berlin).

amplipennis Rehn, 1914 : 165 (*Conchotopoda*). Holotype ♂, CONGO (KINSHASA): Lake Kivu, Idjwi I. (ZMHU, Berlin).

affinis Bolívar, 1922 : 198 (*Peronura*). Holotype ♀, KENYA: Escarpment (MNHN, Paris).

linearis Rehn, 1914 : 167 (*Conchotopoda*). Holotype ♂, UGANDA: Mpororo (ZMHU, Berlin).

? *karschi* Rehn, 1914 : 161 (*Pachypyga*). Holotype ♀, CONGO (KINSHASA): Ruanda, Lake. Mohasi (ZMHU, Berlin).

longipes Bolívar, 1922 : 197 (*Peronura*). Holotype ♀, KENYA: Naivasha (MNHN, Paris).

magna Ragge, 1960 : 304. Holotype ♂, ETHIOPIA: nr. Scillare (BMNH, London).

media Ragge, 1960 : 299. Holotype ♂, ZAMBIA: Lusaka (BMNH, London).

meruensis Sjöstedt, 1909 : 130 (*Plegmatoptera*). Lectotype ♂, TANZANIA: Mt. Meru (NR, Stockholm); designated by Ragge (1960 : 311).

montivaga Sjöstedt, 1909 : 129 (*Peronura*). Lectotype ♂, TANZANIA: Kilimanjaro (NR, Stockholm); designated by Ragge (1960 : 312).

nomima Karsch, 1896 : 327 (*Peronura*). 1 ♂, 2 ♀ syntypes, TANZANIA: Mpwapwa (ZMHU, Berlin).

nuda Ragge, 1960 : 309. Holotype ♂, SUDAN: Imatong Mtns., Kippia (BMNH, London).

regularis Bolívar, 1922 : 200 (*Eupantolepta*). Holotype ♂, KENYA: S. of Lake Rudolph (MNHN, Paris).

? *reticulata* Karsch, 1889 : 429 (*Plegmatoptera*). Holotype ♀, TANZANIA: Lake Tanganyika (ZMHU, Berlin).

- ruspolii*** Schulthess, 1898 : 204 (*Conchotopoda*). Holotype ♂, KENYA: Balessa (MCSN, Genoa).
 ? *rivae* Schulthess, 1898 : 201 (*Peronura*). Holotype ♀, ETHIOPIA: Dolo (MCSN, Genoa).
trochlearis Bolívar, 1922 : 199. Holotype ♂, KENYA: S. of Lake Rudolph (MNHN, Paris).
 ? *viridis* Chopard, in Chopard & Kevan, 1954: 318 (*Peronura*). Holotype ♀, KENYA: Isiolo distr., Bambota (BMNH, London).
samburu Kevan, in Chopard & Kevan, 1954: 320 (*Peronura*). Holotype ♂, KENYA: Samburu (BMNH, London).
- serifera*** Schaum, 1853 : 778. Holotype ♀, MOZAMBIQUE (ZMHU, Berlin).
- vosseleri*** Karny, 1910 : 52 (*Pantolepta*). 10 ♂ syntypes, SOUTH WEST AFRICA: Okahandja;
 1 ♂ syntype, SOUTH WEST AFRICA: Kung-Buschmannland (ZMHU, Berlin).
- multivenosa*** Chopard, 1935 : 2 (*Pantolepta*). Holotype ♂, MOZAMBIQUE: Nova Chupanga, nr. Chemba (MNHN, Paris).
- somali*** Schulthess, 1898 : 200 (*Peronura*). Holotype ♀, SOMALI REPUBLIC: Lugh (MCSN, Genoa).
- peeli*** Burr, 1900 : 44 (*Rhegmatopoda*). Holotype ♂, SOMALI REPUBLIC: Whardi Datal (UM, Oxford).
- stuhlmanni*** Karsch, 1896 : 329 (*Conchotopoda*). Holotype ♂, "AFRICA ORIENTALIS": Buginda (ZMHU, Berlin).
- kasindina*** Rehn, 1914 : 160 (? *Horatosphaga*). Holotype ♀, CONGO (KINSHASA): Kasindi (ZMHU, Berlin).
- stylifera*** Karny, 1910 : 51 (*Pantolepta*). Holotype ♂, SOUTH WEST AFRICA: Windhoek (ZMHU, Berlin).
- vicina*** Chopard, in Chopard & Kevan, 1954 : 318 (*Peronura*). Holotype ♀, KENYA: Marsabit, Chopra Gof (BMNH, London).

ITOKIA Sjöstedt

- Itokiia*** Sjöstedt, 1901 : 10. Type-species: *Itokiia silvarum* Sjöstedt, by monotypy.
- silvarum*** Sjöstedt, 1901 : 10. Holotype ♂, CAMEROUN: Itoki (NR, Stockholm).

IVENSIA Bolívar

- Ivensia*** Bolívar, 1890 : 218. Type-species: *Ivensia uncinata* Bolívar, by monotypy.
- uncinata*** Bolívar, 1890 : 218. Holotype ♂, "Quango" (MLZA, Lisbon).

JAPYGOPHANA Carl

- Japygophana*** Carl, 1921 : 301. Type-species: *Japygophana peloti* Carl, by monotypy.
- peloti*** Carl, 1921 : 302. Holotype ♂, GABON (MHN, Geneva).

KEVANIELLA Chopard

- Kevaniella*** Chopard, in Chopard & Kevan, 1954 : 332. Type-species: *Kevaniella bipunctata* Chopard, by original designation.
- bipunctata*** Chopard, in Chopard & Kevan, 1954: 333. Holotype ♂, KENYA: Mandera distr., Rahmu (BMNH, London).

LAMECOSOMA Ragge

- Lamecosoma*** Ragge, 1960a : 331. Type-species: *Lamecosoma tenuis* Ragge, by original designation.
- inermis*** Ragge, 1961b : 217. Holotype ♂, KENYA: Nairobi, Karura Forest (BMNH, London).
tenuis Ragge, 1960a : 332. Holotype ♂, ZAMBIA: Kipundu (MRAC, Tervuren).

LEIODONTOCERCUS Chopard

Leiodontocercus Chopard, 1954 : 83. Type-species: *Leiodontocercus angustipennis* Chopard, by original designation.

angustipennis Chopard, 1954 : 84. Holotype ♂, GUINEA: Nimba (MNHN, Paris).

condylus Ragge, 1962b : 15. Holotype ♂, CONGO (KINSHASA): Kibali-Ituri, Yindi (MRAC, Tervuren).

malleus Ragge, 1962b : 14. Holotype ♂, GHANA: Western Region, nr. Wiawso (BMNH, London).

LEPTOPHYES Fieber

Leptophyes Fieber, 1853 : 174. Type-species: *Barbitistes albovittatus* Kollar, by monotypy (type-locality: Austria).

antinorii Bormans, 1880 : 217. Holotype ♀, ETHIOPIA: Shoa (MCSN, Genoa).

LETANA Walker

Letana Walker, 1869 : 277. Type-species: *Letana linearis* Walker, by monotypy (type-locality: India).

Pyrrhicia Stål, 1873 : 41. Type-species: *Phaneroptera rubescens* Stål, by original designation (type-locality: Hong Kong).

Pyrrhizia Brunner, 1891 : 55. (Unjustified emendation.)

conradti Bolívar, 1906 : 329 (*Pyrrhicia*). Holotype ♂, CAMEROUN (IEE, Madrid).

zanzibarica Brunner, 1891 : 55 (*Pyrrhizia*). Holotype ♂, "ZANZIBAR" (NM, Vienna).

MANGOMALOBA Sjöstedt

Mangomaloba Sjöstedt, 1902 : 14. Type-species: *Mangomaloba monticola* Sjöstedt, by monotypy.

angustipennis Chopard, 1958 : 76. Holotype ♂, PRINCIPE (MNHN, Paris).

latipennis Chopard, 1954 : 86. Holotype ♂, GUINEA: Nimba (MNHN, Paris).

monticola Sjöstedt, 1901 : 15. Holotype ♂, CAMEROUN: Mt. Cameroon (NR, Stockholm).

royi Chopard, 1954 : 85. Holotype ♂, GUINEA: Nimba (MNHN, Paris).

MELIDIA Stål

Melidia Stål, 1876 : 60. Type-species: *Melidia brunneri* Stål, by monotypy.

brunneri Stål, 1876 : 60. Holotype ♀, SOUTH WEST AFRICA: Damara (NR, Stockholm).

kenyensis Chopard, in Chopard & Kevan, 1954 : 326. Holotype ♀, KENYA: Moyale distr., Yasere (BMNH, London).

laminata Chopard, in Chopard & Kevan, 1954 : 324. Holotype ♂, KENYA: Wajir distr., El Katulo (BMNH, London).

MERUTERRANA Sjöstedt

Meruterrana Sjöstedt, 1912 : 10. Type-species: *Meruterrana elegans* Sjöstedt, by monotypy.

elegans Sjöstedt, 1912 : 12. 4 ♂ syntypes, TANZANIA: Mt. Meru (NR, Stockholm).

MILITITSA Burr

Milititsa Burr, 1900 : 42. Type-species: *Milititsa somaliensis* Burr, by monotypy.

somaliensis Burr, 1900 : 43. Holotype ♀, SOMALI REPUBLIC: Whardi Datal (UM, Oxford).

MILTINOBATES Sjöstedt

Miltinobates Sjöstedt, 1901 : 15. Type-species: *Miltinobates blandus* Sjöstedt, by monotypy.

blandus Sjöstedt, 1901 : 16. Holotype ♂, "KONGO" (NR, Stockholm).

MONTEIROA Karsch

Monteiroa Karsch, 1889 : 458. Type-species: *Monteiroa latifrons* Karsch, by monotypy.

latifrons Karsch, 1889 : 458. 1 ♂, 1 ♀ syntypes, MOZAMBIQUE: Delagoa Bay (ZMHU, Berlin).

MONTICOLARIA Sjöstedt

Monticolaria Sjöstedt, 1909 : 128. Type-species: *Monticolaria meruensis* Sjöstedt, by PRESENT DESIGNATION.

Both the species listed below were originally included in *Monticolaria* Sjöstedt, but neither was designated as the type-species. I here designate *Monticolaria meruensis* Sjöstedt as the type-species of *Monticolaria* Sjöstedt.

kilimandjarica Sjöstedt, 1909 : 129. 2 ♂ syntypes, TANZANIA: Kilimanjaro (NR, Stockholm).
meruensis Sjöstedt, 1909 : 128. 10 ♂, 3 ♀ syntypes, TANZANIA: Meru (NR, Stockholm).

MORGENIA Karsch

Morgenia Karsch, 1890b : 261. Type-species: *Morgenia hamuligera* Karsch, by monotypy.

hamuligera Karsch, 1890b : 263. Holotype ♂, CAMEROUN: Kribi (ZMHU, Berlin).

metica Karsch, 1893 : 196. Holotype ♂, CAMEROUN: Victoria (ZMHU, Berlin).

modulata Karsch, 1896 : 340. 1 ♂, 1 ♀ syntypes, CAMEROUN: Lolodorf (ZMHU, Berlin).

rubicornis Sjöstedt, 1913 : 4. Holotype ♂, CONGO (KINSHASA): Mukimbungu (NR, Stockholm).

spathulifera Griffini, 1908 : 209. 1 ♂ syntype, "Bussanga"; 1 ♀ syntype, CAMEROUN: Mukonje Farm (IRSNB, Brussels).

MYLLOCENTRUM Ragge

Mylocentrum Ragge, 1962b : 15. Type-species: *Phlaurocentrum stigmosum* Karsch, by original designation.

stigmosum Karsch, 1896 : 336 (*Phlaurocentrum*). Holotype ♀, CAMEROUN: Lolodorf (ZMHU, Berlin).

ODONTURA Rambur

Odontura Rambur, 1838 : 44. Type-species: *Barbitistes glabricula* Charpentier, by subsequent designation (Kirby, 1906 : 386) (type-locality: Portugal).

algerica Brunner, 1878 : 75. 1 ♂, 1 ♀ syntypes, ALGERIA (NM, Vienna).

borrei Bolívar, 1878 : 71. Unknown number of ♀ syntypes, ALGERIA: Constantine Province (IRSNB, Brussels).

- brevis* Werner, 1932 : 114. Holotype ♂, MOROCCO: Ifrane (NM, Vienna).
- capensis* Walker, 1869 : 242. Holotype ♀, SOUTH AFRICA (BMNH, London).
- liouvillei* Werner, 1929 : 184. Unknown number of ♂ syntypes, MOROCCO: nr. Tiflet, Mamora Forest; 1 ♂ syntype, MOROCCO: Malay Idris (NM, Vienna).
- maroccana* Bolívar, 1908 : 333 (as var. of *Odontura spinulicauda*; raised to sp. by Bolívar, 1914 : 237). 7 ♂, 1 ♀ syntypes, MOROCCO: Marrakesh (IEE, Madrid).
- technicus* Morales, 1947 : 250 (as subsp. of *Odontura liouvillei*). Holotype ♂, IFNI: Zoco Tlata Sboya (IEE, Madrid).
- microptera* Chopard, 1943 : 107. Holotype ♀, ALGERIA: Ain Mlila (MNHN, Paris).
- moghrebica* Morales, 1950 : 163. Holotype ♂, MOROCCO: Isaguen (IEE, Madrid).
- plasconi* Ebner, 1915 : 419. 3 ♂, 1 ♀ syntypes, TANZANIA: Tassamaganga (NM, Vienna).
- pulchra* Bolívar, 1914 : 237. 3 ♂, 3 ♀ syntypes, MOROCCO: Tangier (IEE, Madrid).
- quadridentata* Krauss, 1893 : xcvi. Unknown number of syntypes of both sexes, ALGERIA: Tessala (type-depository unknown).
- terniensis* Finot, 1893 : xxx. Unknown number of syntypes of both sexes, ALGERIA: Terni, Sebdou and Tlemcen (MNHN, Paris).
- spinulicauda* Rambur, 1838 : 45. Lectotype ♂, SPAIN (IEE, Madrid); designated by Uvarov (1948 : 380).
- stenoxypha* Fieber, 1853 : 187 (*Barbitistes*). Holotype ♀, SICILY (type-depository unknown).
- uvarovi* Werner, 1929 : 184. Holotype ♂, MOROCCO: Fez (NM, Vienna).

OXYECOUS Chopard

- Oxyecous* Chopard, 1935 : 4. Type-species: *Oxyecous lesnei* Chopard, by monotypy.
- apertus* Ragge, 1956b : 187. Holotype ♂, KENYA: Rabai (BMNH, London).
- lesnei* Chopard, 1935 : 4. Holotype ♂, MOZAMBIQUE: nr. Vile Pery (MNHN, Paris).
- magnus* Ragge, 1956b : 188. Holotype ♂, TANZANIA: Kilosa (BMNH, London).
- undulatus* Ragge, 1956b : 190. Holotype ♂, UGANDA: Bugoma Forest (BMNH, London).

PARAPYRRHICIA Brunner

- Parapyrrhicia* Brunner, 1891 : 149. Type-species: *Parapyrrhicia zanzibarica* Brunner, by monotypy.
- zanzibarica* Brunner, 1891 : 149. Holotype ♀, "ZANZIBAR" (NM, Vienna).

PARDALOTA Brunner

- Pardalota* Brunner, 1878 : 133. Type-species: *Pardalota versicolor* Brunner, by monotypy.
- asymmetrica* Karsch, 1896 : 330. 7 syntypes of both sexes, "AFRICA ORIENTALIS CENTRALIS": Kirewia; Itimba; Bundeko; Bugundi; Sossian-Ngoroine, Massai Territory; W. Victoria Nyansa, Ussoga-Uganda (ZMHU, Berlin).
- cloetensi* Griffini, 1908a : 77. Holotype ♀, CONGO (KINSHASA): Beni Bendi, Sankuru (IRSNB, Brussels).
- haasi* Griffini, 1908a : 76. 2 ♂, 1 ♀ syntypes, CONGO (KINSHASA): Luluabourg (IRSNB, Brussels).
- karschiana* Enderlein, 1907 : 197. 2 ♂ syntypes, TANZANIA: Dar es Salaam (1 in ZMHU, Berlin; 1 in IZPAN, Warsaw).
- reimeri* La Baume, 1911 : 317. 2 ♀, 3 nymphal syntypes, TANZANIA: Lindi (ZMHU, Berlin).
- superba* Sjöstedt, 1913 : 2. Holotype ♂, TANZANIA: Lukuledi (NR, Stockholm).
- versicolor* Brunner, 1878 : 134. Unknown number of syntypes of both sexes, "Chinchoxa in Benguela" (1 ♂ in ZMHU, Berlin; 1 ♂ in NM, Vienna).

PERONURA Karsch

Peronura Karsch, 1889 : 426. Type-species: *Peronura clavigera* Karsch, by subsequent designation (Kirby, 1906 : 391).

clavigera Karsch, 1889 : 427. 2 ♀ syntypes, KENYA: Mombasa (ZMHU, Berlin).

PEROPYRRHICIA Brunner

Peropyrrhicia Brunner, 1891 : 37. Type-species: *Dichopetala massaiae* Bormans, by monotypy.

cooperi Uvarov, 1934 : 598. Holotype ♂, ETHIOPIA: Jem-Jem Forest (BMNH, London).

maculata Schulthess, 1898 : 198. 1 ♂, 1 ♀ syntypes, ETHIOPIA: Biddwara (MCSN, Genoa).

massaiae Bormans, 1880 : 218 (*Dichopetala*). 2 ♂ syntypes (1 nymphal), ETHIOPIA: Let-Marefia (MCSN, Genoa).

scotti Uvarov, 1934 : 597. Holotype ♂, ETHIOPIA: between Jem-Jem and Addis Ababa (BMNH, London).

PHANEROPTERA Serville

Phaneroptera Serville, 1831 : 158. Type-species: *Gryllus falcata* Poda, by subsequent designation (Hemming, 1944 : 211) (type-locality unknown).

Anerota Caudell, 1921 : 488. Type-species: *Gryllus falcata* Poda, by original designation.

Paranerota Karny, 1926 : 105. Type-species: *Phaneroptera gracilis* Burmeister, by original designation.

Euanerota Karny, 1927 : 12. Type-species: *Phaneroptera brevis* Serville, by tentative original designation (see Ragge, 1956 : 206) (type-locality: Java).

acaciae Chopard, in Chopard & Kevan, 1954 : 323. Holotype ♂, KENYA: Mandera distr., Rahmu (BMNH, London).

africana Steinmann, 1966 : 411. Holotype ♂, GUINEA: Conakry (TM, Budapest).

albida Walker, 1869 : 335. Holotype ♂, no data (BMNH, London).

fragilis Ragge, 1960b : 245. Holotype ♂, ETHIOPIA: Danot (BMNH, London).

gracilis Burmeister, 1838 : 690. 1 ♂, 2 ♀ syntypes, JAVA (type-depository unknown).

roseata Walker, 1869 : 343. Holotype ♀, INDIA (BMNH, London).

marginalis Brunner, 1878 : 214. Holotype ♀, SOUTH AFRICA (ZMHU, Berlin).

indica Brunner, 1878 : 215. Lectotype ♂, INDIA: Himalayas (NM, Vienna); designated by Ragge (1960b : 250).

elongata Brunner, 1878 : 217. Lectotype ♀, JAVA (ZMHU, Berlin); designated by Ragge (1956a : 251).

subcarinata Bolívar, 1900 : 764. Holotype ♂, INDIA: Madura (IEE, Madrid).

longispina Ragge, 1956a : 244. Holotype ♂, UGANDA: Ruwenzori Range, Kilembe (BMNH, London).

maculosa Ragge, 1956a : 243. Holotype ♂, CAMEROUN: Lolodorf (ZMHU, Berlin).

magna Ragge, 1956a : 242. Holotype ♂, UGANDA: Mabira Forest (BMNH, London).

minima Brunner, 1878 : 214. Holotype ♂, EGYPT (ZMHU, Berlin).

nana nana Fieber, 1853 : 173 (as sp.). Type-locality: PORTUGAL; type-material lost.

quadripunctata Brunner, 1878 : 212. Lectotype ♂, ITALY: Trieste (NM, Vienna); designated by Ragge (1956 : 233).

nana sparsa Stål, 1856 : 170 (as sp.). Holotype ♀, SOUTH AFRICA: "Caffraria" (NR, Stockholm).

lurida Walker, 1869 : 339. Holotype ♂, "W. AFRICA" (erroneously cited as "Natal" in original description) (BMNH, London).

tetrasticta Gerstaecker, 1869 : 215. Holotype ♂, TANZANIA: Uru (ZMHU, Berlin).

- conspersa* Stål, 1874 : 29 (apparently proposed unnecessarily as replacement name for *Phaneroptera sparsa* Stål).
punctulata Burr, 1900 : 41. Holotype ♂, SOMALI REPUBLIC: Whardi Datal (UM, Oxford).
tenuicerca Ramme, 1951 : 348. Holotype ♂, LEBANON: Djezin (ZMHU, Berlin).
nigropunctata Chopard, 1955 : 269. Holotype ♂, SOUTH AFRICA: Cape Province, Drakensbergen, 8 miles E.N.E. of Rhodes (UZI, Lund).
parva Ragge, 1956a : 240. Holotype ♂, CONGO (KINSHASA): Kasongo (MRAC, Tervuren).

PHANEROPTILA Uvarov

Phaneroptila Uvarov, in Uvarov & Popov, 1957 : 363. Type-species: *Phaneroptila insularis* Uvarov, by monotypy.

insularis Uvarov, in Uvarov & Popov, 1957 : 363. Holotype ♂, SOCOTRA: Hijama (BMNH, London).

PHAULULA Bolívar

Phaula Brunner, 1878 : 167. Type-species: *Phaula laevis* Brunner, by subsequent designation (Kirby, 1906 : 425) (type-locality: Philippine Is.). (Homonym of *Phaula* Thomson, 1857 : 303.)

Phaulula Bolívar, 1906 : 347 (proposed as replacement name for *Phaula* Brunner).

inconspicua Brunner, 1891 : 83 (*Phaula*). Holotype ♀, CAMEROUN (NM, Vienna).

lenzi Brunner, 1891 : 83 (*Phaula*). Unknown number of ♂ syntypes, CAMEROUN (Lübeck, museum unknown) and GABON (NM, Vienna).

PHLAUROCENTRUM Karsch

Phlaurocentrum Karsch, 1889 : 445. Type-species: *Phlaurocentrum latevittatum* Karsch, by monotypy.

latevittatum Karsch, 1889 : 446. Holotype ♀, CONGO (KINSHASA): Kuako to Kimpoko (ZMHU, Berlin).

lobatum Ragge, 1962b : 9. Holotype ♂, CONGO (KINSHASA): Eala (MRAC, Tervuren).

maculatum Ragge, 1962b : 10. Holotype ♂, CONGO (KINSHASA): Mawambi-Irumu (NM, Vienna).

mecopodoides Karsch, 1892a : 321. Holotype ♂, CAMEROUN: Barombi (ZMHU, Berlin).

tuberosum Ragge, 1962b : 8. Holotype ♂, UGANDA: Mabira Forest (BMNH, London).

turbatum Walker, 1869 : 340 (*Phaneroptera*). Holotype ♂, "CONGO" (BMNH, London).

PHYSOCORYPHA Karsch

Physocorypha Karsch, 1896 : 344. Type-species: *Physocorypha politurata* Karsch, by monotypy.

politurata Karsch, 1896 : 345. 1 ♀ syntype, CAMEROUN: Victoria; 1 ♀ syntype, CAMEROUN: Mundame (ZMHU, Berlin).

PLANGIA Stål

Plangia Stål, 1873 : 40. Type-species: *Phylloptera graminea* Serville, by original designation.

graminea Serville, 1839 : 405 (*Phylloptera*). Unknown number of syntypes of both sexes, SOUTH AFRICA: Cape of Good Hope (lost).

natalensis Walker, 1869 : 379 (*Phylloptera*). Holotype ♀, SOUTH AFRICA: Natal (BMNH, London).

- karschi*** Chopard, 1954 : 44. Holotype ♀, GUINEA: Nimba, Nion (MNHN, Paris).
laminifera Karsch, 1896 : 343. Holotype ♀, CAMEROUN: Lolodorf (ZMHU, Berlin).
nebulosa Karsch, 1890c : 366. Holotype ♀, CAMEROUN: Barombi (ZMHU, Berlin).
unimaculata Chopard, 1955 : 271. Holotype ♂, SOUTH AFRICA: Cape Province, Tzitzikama Forest, Storms River Mouth (UZI, Lund).
villiersi Chopard, 1954 : 45. Holotype ♀, GUINEA: Nimba (MNHN, Paris).

PLANGIOIDES Chopard

Plangiodes Chopard, in Chopard & Kevan, 1954 : 330. Type-species: *Plangiodes carinatus* Chopard, by original designation.

carinatus Chopard, in Chopard & Kevan, 1954 : 330. Holotype ♂, KENYA: Garissa distr., Afwein (BMNH, London).

PLANGIOLA Bolívar

Plangiola Bolívar, 1906 : 336. Type-species: *Plangiola herbacea* Bolívar, by monotypy.

herbacea Bolívar, 1906 : 337. 2 ♂, 1 ♀ syntypes, CAMEROUN (IEE, Madrid).

PLANGIOPSIS Karsch

Plangiopsis Karsch, 1889 : 459. Type-species: *Plangiopsis semiconchata* Karsch, by monotypy.

adeps Karsch, 1896 : 338. 2 ♀ syntypes, CAMEROUN: Lolodorf (ZMHU, Berlin).

foraminata Karsch, 1892a : 324. Holotype ♂, CAMEROUN: Barombi (ZMHU, Berlin).

schoutedeni Griffini, 1908 : 218. Holotype ♀, CAMEROUN: Mukonje Farm (IRSNB, Brussels).

semiconchata Karsch, 1889 : 460. Holotype ♀, CAMEROUN: Barombi (ZMHU, Berlin).

POECILOGRAMMA Karsch

Poecilogramma Karsch, 1887b : 52. Type-species: *Poecilogramma striatifemur* Karsch, by subsequent designation (Kirby, 1906 : 405).

annulifemur Karsch, 1887b : 53. 2 ♂, 3 ♀ syntypes, TANZANIA: Kawende (ZMHU, Berlin).
striatifemur Karsch, 1887b : 53. 2 ♂ syntypes, KENYA: Mombasa (ZMHU, Berlin).

POREUOMENA Brunner

Poreuomena Brunner, 1878 : 187. Type-species: *Poreuomena africana* Brunner, by monotypy.

africana Brunner, 1878 : 187. Holotype ♂, GABON (MHN, Geneva).

duponti Griffini, 1908a : 84. 1 ♂, 6 ♀ syntypes, CAMEROUN: Mukonje Farm (IRSNB, Brussels).

forcipata Sjöstedt, 1901 : 12. Holotype ♂, CAMEROUN (NR, Stockholm).

gladiator Bolívar, 1906 : 337. Holotype ♂, CAMEROUN (IEE, Madrid).

lamottei Chopard, 1954 : 40. 1 ♂ syntype, GUINEA: Nimba, N'Zo; 1 ♀ syntype, GUINEA: Nimba, Yalanzou (MNHN, Paris).

wilverthi Griffini, 1908a : 85. Holotype ♂, CONGO (KINSHASA): Umangi (IRSNB, Brussels).

PREUSSIA Karsch

Preussia Karsch, 1890c : 364. Type-species: *Preussia lobatipes* Karsch, by monotypy.

lobatipes Karsch, 1890c : 365. Holotype ♀, CAMEROUN: Barombi (ZMHU, Berlin).

PRONOMAPYGA Rehn

Pronomapyga Rehn, 1914 : 171. Type-species: *Pronomapyga grandis* Rehn, by original designation.

grandis Rehn, 1914 : 172. Holotype ♂, "Ost-AFRIKA" (ZMHU, Berlin).
graueri Rehn, 1914 : 171. Holotype ♀, "Ost-AFRIKA" (ZMHU, Berlin).

PROSPHAGA Ragge

Prosphega Ragge, 1960a : 322. Type-species: *Pantolepta calaharica* Karny, by original designation.

calaharica Karny, 1910 : 52 (*Pantolepta*). Holotype ♂, SOUTH AFRICA: Bechuanaland, Kalahari, Ku Gudie, nr. Pitsani (ZMHU, Berlin).

splendens Ragge, 1960a : 324. Holotype ♂, ETHIOPIA: Wardere (BMNH, London).

PSEUDOPYRRHIZIA Brunner

Pseudopyrrhizia Brunner, 1891 : 110. Type-species: *Pseudopyrrhizia punctata* Brunner, by monotypy.

punctata Brunner, 1891 : 110. Holotype ♂, "ZANZIBAR" (NM, Vienna).

SCHUBOTZACRIS Rehn

Schubotzcris Rehn, 1914 : 169. Type-species: *Schubotzcris producta* Rehn, by original designation.

producta Rehn, 1914 : 169. Holotype ♂, CONGO (KINSHASA): Mboga (ZMHU, Berlin).

STENAMBLYPHYLLUM Karsch

Stenamblyphyllum Karsch, 1896 : 326. Type-species: *Stenamblyphyllum dilutum* Karsch, by monotypy.

dilutum Karsch, 1896 : 327. Lectotype ♀, CAMEROUN: Victoria (ZMHU, Berlin); designated by Ragge (1962a : 309).

SYMMETROPLEURA Brunner

Symmetropleura Brunner, 1878 : 245. Type-species: *Symmetropleura laevicauda* Brunner, by subsequent designation (Kirby, 1906 : 446) (type-locality: South America, " Bahia ").

Cameronia Karsch, 1889 : 450. Type-species: *Symmetropleura africana* Brunner, by monotypy. (See p. 77.)

africana Brunner, 1878 : 246. Holotype ♂, "CONGO" (NM, Vienna).

plana Walker, 1869 : 339 (*Phaneroptera*). Holotype ♂, SOUTH AFRICA: Natal (BMNH, London).

TAPIENA Bolívar

Tapeina Brunner, 1878 : 163. Type-species: *Tapeina acutangula* Brunner, by monotypy (type-locality : India). (Homonym of *Tapeina* Le Peletier & Serville, 1828 : 545.)

Tapiena Bolívar, 1906 : 334 (proposed as replacement name for *Tapeina* Brunner).

minor Bolívar, 1906 : 334. Holotype ♀, GHANA: " Ashante " (IEE, Madrid).

TERPNISTRIA Stål

Terpnistria Stål, 1873 : 42. Type-species: *Phaneroptera zebra* Serville, by original designation.

lobulata Stål, 1876 : 61. Holotype ♀, SOUTH WEST AFRICA: Ovambo (NR, Stockholm) (the additional citation of "Caffraria" is probably erroneous).

tuberculata Chopard, in Chopard & Kevan, 1954 : 326. Holotype ♂, KENYA: Northern Frontier Distr., Garissa (BMNH, London).

zebrata Serville, 1839 : 424 (*Phaneroptera*). Holotype ♂, SOUTH AFRICA: Cape of Good Hope (lost).

TETRACONCHA Karsch

Tetraconcha Karsch, 1890a : 61. Type-species: *Tetraconcha fenestrata* Karsch, by monotypy. *Tellidia* Bolívar, 1893 : 177. Type-species: *Tellidia longipes* Bolívar, by monotypy.

banzyvilleana Griffini, 1909 : 11. Holotype ♀, CONGO (KINSHASA): Banzyville, Jusoro (MRAC, Tervuren).

fenestrata Karsch, 1890a : 62. Holotype ♂, CAMEROUN (ZMHU, Berlin).

longipes Bolívar, 1893 : 178 (*Tellidia*). Holotype ♀, IVORY COAST: Assinie (IEE, Madrid).

smaragdina Brunner, 1891 : 116. Holotype ♂, CAMEROUN (Lübeck, museum unknown).

stichyrata Karsch, 1890c : 360. Holotype ♂, CAMEROUN: Barombi (ZMHU, Berlin).

scalaris Brunner, 1891 : 116. Holotype ♂, GABON (NM, Vienna).

TRIGONOCORYPHA Stål

Trigonocorypha Stål, 1873 : 39. Type-species: *Locusta crenulata* Thunberg (= *Trigonocorypha unicolor* (Stoll)), by original designation (type-locality unknown).

tihamae Uvarov, 1952 : 177. Holotype ♂, SAUDI ARABIA: near Qunfida, Wadi Qanuna (BMNH, London).

TROPIDONOTACRIS Chopard

Tropidonotacris Chopard, in Chopard & Kevan, 1954 : 321. Type-species: *Tropidonotacris carinata* Chopard, by original designation.

amabilis Ragge, 1957 : 121. Holotype ♂, ETHIOPIA: Ual Ual (BMNH, London).

carinata Chopard, in Chopard & Kevan, 1954 : 322 (as *Tropidonotacris carinatus*; corrected by Ragge, 1957 : 119). Holotype ♂, KENYA: Mandera distr., Takabba (BMNH, London).

grandis Ragge, 1957 : 122. Holotype ♂, TANZANIA: Little Mahenge (BMNH, London).

TROPIDOPHRYS Karsch

Tropidophrys Karsch, 1896 : 340. Type-species: *Tropidophrys amydra* Karsch, by monotypy.

amydra Karsch, 1896 : 341. Holotype ♂, CAMEROUN: Victoria (ZMHU, Berlin).

TYLOPSIS Fieber

Centrophorus Fischer, 1846 : 361. Type-species: *Centrophorus spinosus* Fischer (= *Tylopsis lilifolia* (Fabricius)), by monotypy. (Homonym of *Centrophorus* Müller & Henle, 1837 : 115.) *Tylopsis* Fieber, 1853 : 172. Type-species: *Locusta lilifolia* Fabricius, by monotypy.

- ampla** Ragge, 1964 : 315. Holotype ♂, ANGOLA: Moxico distr., upper Mu-Simoj R. (BMNH, London).
- bilineolata** Serville, [1838]: 419 (*Phaneroptera*). Holotype ♂, SOUTH AFRICA: Cape of Good Hope (lost).
- attenuata* Walker, 1869 : 338 (*Phaneroptera*). Holotype ♀, SOUTH AFRICA (BMNH, London).
- marginata* Brunner, 1891 : 113. Holotype ♀, SOUTH AFRICA: Natal, Durban (NM, Vienna).
- brevis** Ragge, 1964 : 314. Holotype ♂, CONGO (KINSHASA): Volcan Nyamлагира (MRAC, Tervuren).
- continua** Walker, 1869 : 337 (*Phaneroptera*). Lectotype ♂, SOUTH AFRICA: Natal, Durban (BMNH, London); designated by Ragge (1964 : 318).
- vicaria** Walker, 1869 : 338 (*Phaneroptera*). Holotype ♂, SOUTH AFRICA: Natal, Durban (BMNH, London).
- longipennis* Stål, 1876 : 58. Holotype ♂, SOUTH WEST AFRICA: Damara (NR, Stockholm).
- vittata* Brunner, 1878 : 229. Lectotype ♂, SOUTH AFRICA: Natal, Durban (NM, Vienna); designated by Ragge (1964 : 318).
- inhamata* Karsch, 1889 : 453. Lectotype ♂, MOZAMBIQUE: Delagoa Bay (ZMHU, Berlin).
- dispar** Sjöstedt, 1909 : 135. Lectotype ♂, TANZANIA: Mt. Meru (NR, Stockholm); designated by Ragge (1964 : 309).
- dubia** Giglio-Tos, 1907 : 3. 1 ♂, 1 nymphal syntype, UGANDA: Ibanda; 1 ♀ syntype, UGANDA: Mitiana (IMZU, Turin).
- fissa** Ragge, 1964 : 312. Holotype ♂, UGANDA: Entebbe (BMNH, London).
- gracilis** Chopard, in Chopard & Kevan, 1954 : 328. Holotype ♂, KENYA: Mandera distr., Damassa (BMNH, London).
- irregularis** Karsch, 1893 : 130. Lectotype ♂, TOGO: Bismarckburg (ZMHU, Berlin); designated by Ragge (1964 : 307).
- perpulchra* Burr, 1900 : 43. Holotype ♂, SOMALI REPUBLIC: Whardi Datal (UM, Oxford).
- obscuripes* Chopard, 1945 : 166. Lectotype ♂, CAMEROUN: Bambouto Mtns. (MNHN, Paris); designated by Ragge (1964 : 307).
- villiersi* Chopard, 1950 : 133. Lectotype ♂, NIGER: Agadès (MNHN, Paris); designated by Ragge (1964 : 307).
- lamottei* Chopard, 1954 : 42. Lectotype ♂, GUINEA: Nimba, Serengbara (MNHN, Paris); designated by Ragge (1964 : 307).
- liliifolia** Fabricius, 1793 : 36 (*Locusta*). Type-locality: ITALY; type-material lost.
- gracilis* Germar, 1817 : 251 (*Locusta*). Holotype ♀, YUGOSLAVIA: Zadar (lost).
- liliifolia* Rambur, 1838 : 44 (*Locusta*). (Unjustified emendation.)
- praeusta* Fischer, 1846 : 142 (*Phaneroptera*). Holotype ♀, U.S.S.R.: Azerbaijan, Kirovabad (lost).
- spinosus* Fischer, 1846 : 362 (*Centrophorus*). Unknown number of nymphal syntypes of both sexes, U.S.S.R.: Crimea (lost).
- margineguttata* Serville, 1839 : 442 (*Phaneroptera*). 1 ♂ syntype, SARDINIA; 1 ♂ syntype, SICILY (lost).
- ? *coi* Jannone, 1936 : 147. Holotype ♂. DODECANESE: Kos (ILEA, Portici).
- rubrescens** Kirby, 1900 : 216. Holotype ♂, MALAWI: Zomba (BMNH, London).
- punctulata* Kirby, 1900 : 216. Lectotype ♂, MALAWI: Zomba (BMNH, London); designated by Ragge (1964 : 320).
- meruensis* Sjöstedt, 1909 : 134. Lectotype ♂, TANZANIA: Mt. Meru (NR, Stockholm); designated by Ragge (1964 : 320).
- confluens* Karny, 1915 : 124. Holotype ♀, GUINEA: Upper Niger (NM, Vienna).

VOSSIA Brunner

Vossia Brunner, 1891 : 139. Type-species: *Vossia obesa* Brunner, by monotypy.

obesa Brunner, 1891 : 140. Holotype ♀, CAMEROUN (NM, Vienna).

WEISSENBORНИA Karsch

Weissenbornia Karsch, 1888 : 65. Type-species: *Weissenbornia praestantissima* Karsch, by monotypy.

praestantissima Karsch, 1888 : 66. Holotype ♂, CAMEROUN: Lowry-Criby-Mündung (ZMHU, Berlin).

ZEUNERIA Karsch

Zeuneria Karsch, 1889 : 443. Type-species: *Zeuneria melanopeza* Karsch, by monotypy.

biramosa Sjöstedt, 1929 : 40. Holotype ♂, CONGO (KINSHASA): Bas-Uele (MRAC, Tervuren).

centralis Rehn, 1914 : 179. Holotype ♂, CONGO (KINSHASA): Ruwenzori (ZMHU, Berlin).

longicercus Sjöstedt, 1929 : 39. Holotype ♂, CONGO (KINSHASA): Inkisi (MRAC, Tervuren).

melanopeza Karsch, 1889 : 443. Holotype ♀, CAMEROUN: Barombi (ZMHU, Berlin).

REFERENCES

- ANDER, K. 1939. Vergleichend-anatomische und phylogenetische Studien über die Ensifera (Saltatoria). *Opusc. ent. Suppl.* II. viii + 306 pp., 172 figs.
- BEI-BIENKO, G. YA. 1954. Orthoptera 2 (2). Phaneropterinae. [In Russian.] *Fauna SSSR* 59, 384 pp., 210 figs.
- 1964. *Key to the insects of the European part of the U.S.S.R. 1. Lower Palaeoptera, with imperfect metamorphosis.* [In Russian.] 936 pp., 344 figs. Moscow.
- BEIER, M. 1955. *Bronn's Kl. Ordn. Tierreichs. Fünfter Band: Arthropoda. III. Abteilung: Insecta. 6. Buch: Embioidea und Orthopteroidea.* i + 304 pp., 197 figs. Leipzig.
- BLATCHLEY, W. S. 1920. *Orthoptera of northeastern America with especial reference to the faunas of Indiana and Florida.* 784 pp., 246 figs., 1 pl. Indianapolis.
- BOLÍVAR, I. 1890. Ortópteros de África del Museo de Lisboa (conclusion). *Jorn. Sci. math. phys. nat.* (2) 1 : 211–232, 1 pl.
- 1893. Voyage de M. Ch. Alluaud dans le territoire d'Assinie (Afrique occidentale) en juillet et août 1886. 14^e Mémoire. Orthoptères. *Annls Soc. ent. Fr.* 62 : 169–184, 1 pl.
- 1900. Les Orthoptères de St-Joseph's College à Trichinopoly (Sud de l'Inde). Pt. II. *Annls Soc. ent. Fr.* 68 : 761–812, 2 pls.
- 1906. Fasgonurídos de la Guinea española. *Mems R. Soc. esp. Hist. nat.* 1 : 327–377, 1 pl.
- 1908. Algunos ortópteros nuevos de España, Marruecos y Canarias. *Boln R. Soc. esp. Hist. nat.* 8 : 317–334.
- 1914. Dermápteros y ortópteros de Marruecos. *Mems R. Soc. esp. Hist. nat.* 8 : 157–238.
- 1922. *Voyage de M. le Baron Maurice de Rothschild en Éthiopie et en Afrique Orientale Anglaise (1904–1905). Résultats scientifiques. Animaux articulés. Première Partie. Orthoptères.* Pp. 169–219, 4 pls. Paris.
- BORMANS, A. DE. 1880. Spedizione Italiana nell'Africa equatoriale. Risultati zoologici. Ortotteri. *Annali Mus. civ. Stor. nat. Genova* 16 : 205–221, 4 figs.
- BRUES, C. T., MELANDER, A. L., & CARPENTER, F. M. 1954. Classification of insects. *Bull. Mus. comp. Zool. Harv.* 108, v + 917 pp., 1219 figs.
- BRUNER, L. 1920. Orthoptera from Africa, being a report upon some Saltatoria mainly from Cameroon contained in the Carnegie Museum. *Ann. Carneg. Mus.* 13 : 92–142.
- BRUNNER VON WATTENWYL, C. 1878. *Monographie der Phaneropteriden.* 401 pp., 8 pls. Wien.
- 1883. Ueber hypertelische Nachahmungen bei Orthopteren. *Verh. zool.-bot. Ges. Wien* 33 : 247–249, 1 pl.

- BRUNNER VON WATTENWYL, C. 1891. Additamenta zur Monographie der Phaneropteriden. *Verh. zool.-bot. Ges. Wien* **41** : 1-196, 2 pls.
- 1893. Révision du système des Orthoptères et description des espèces rapportées par M. Leonardo Fea de Birmanie. *Annali Mus. civ. Stor. nat. Genova* (2) **13** : 5-230, 6 pls.
- BURMEISTER, H. 1834-1839. *Handbuch der Entomologie* **2**, xii + 1050 pp. Berlin.
- BURR, M. 1900. On a collection of insects and Arachnids made in 1895 and 1897, by Mr. C. V. A. Peel, F.Z.S., in Somaliland, with descriptions of new species. Pt. 7. Orthoptera. *Proc. zool. Soc. Lond.* **1900** : 35-46, 1 pl.
- 1910. *A synopsis of the Orthoptera of western Europe*. 160 pp. London.
- CARL, J. 1921. Phasgonurides nouveaux du Muséum de Genève. *Rev. suisse Zool.* **28** : 301-309, 5 figs.
- CAUDELL, A. N. 1921. On the orthopterous group Phaneropterae (= Scudderiae), with descriptions of a new genus and species. *J. Wash. Acad. Sci.* **11** : 487-493, 1 fig.
- CHOPARD, L. 1922. Orthoptères et Dermoptères. *Faune Fr.* **3**, 22 pp., 466 figs.
- 1935. Contributions à l'étude de la faune du Mozambique. Voyage de M. P. Lesne (1928-1929). 19. Orthoptères Ensifères. *Mems Estud. Mus. zool. Univ. Coimbra* (I) **85** : 1-32, 18 figs.
- 1938. Mission scientifique de l'Omo. Tome IV. Fasc. 33. Orthoptera. 1. Dictyoptera, Phasmodaea, Ensifera. *Mém. Mus. natn. Hist. nat., Paris* **8** : 89-134, 34 figs.
- 1943. Orthoptéroïdes de l'Afrique du Nord. *Faune Emp. fr.* **1**, 450 pp., 658 figs.
- 1945. Orthoptéroïdes recueillis dans les montagnes du Cameroun par la mission Lepesme, Paulian, Villiers. *Revue fr. Ent.* **11** : 156-178, 27 figs.
- 1947. *Atlas des Aptérygotes et Orthoptéroïdes de France*. 111 pp., 12 figs., 12 pls. Paris.
- 1949. In Grassé, P.-P., [Ed.], *Traité de Zoologie* **9**, 1117 pp., 752 figs., 2 pls. Paris.
- 1950. Contribution à l'étude de l'Air (Mission L. Chopard et A. Villiers). Orthoptéroïdes. *Mém. Inst. fr. Afr. noire* **10** : 127-145, 2 figs.
- 1954. La réserve naturelle intégrale du Mont Nimba. Fasc. II. Pt. III. Orthoptères Ensifères. *Mém. Inst. fr. Afr. noire* **40** (2) : 25-97, 42 figs.
- 1955. In Hanström, B., Brinck, P. & Rudebeck, G., [Edd.], *South African Animal Life* **2**, pp. 266-300 (Orthoptera Ensifera), 26 figs. Stockholm.
- 1958. Mission du Muséum dans les îles du Golfe de Guinée. Entomologie, VI. Orthoptéroïdes. *Bull. Soc. ent. Fr.* **63** : 73-85, 2 figs.
- CHOPARD, L., & KEVAN, D. K. McE. 1954. Orthoptera-Ensifera from Northern Kenya and Jubaland. *Trans. R. ent. Soc. Lond.* **105** : 315-353, 22 figs.
- DIRSH, V. M. 1961. A preliminary revision of the families and subfamilies of Acridoidea (Orthoptera, Insecta). *Bull. Br. Mus. nat. Hist. (Ent.)* **10** : 351-419, 34 figs.
- EBNER, R. 1915. Neue tropische Odonturen (Orthoptera). *Zool. Anz.* **45** : 419-422.
- 1943. Einige Orthoptera Saltatoria von Fernando Poo (Spanisch-Guinea). 28. Beitrag zu den wissenschaftlichen Ergebnissen der Westafrika-Expedition Eidmann 1939/40. *Zool. Anz.* **143** : 259-274, 12 figs.
- ENDERLEIN, G. 1907. *Pardalota karschiana*, eine neue ost-afrikanische Orthoptere. *Zool. Jb. (Syst. Geogr. Biol.)* **25** : 197-200, 1 fig., 1 pl.
- ESSIG, E. O. 1947. *College entomology*. vii + 900 pp., 308 figs. New York.
- FABRICIUS, J. C. 1793. *Entomologia systematica emendata et aucta* **2**, viii + 519 pp. Copenhagen.
- FIEBER, F. X. 1853. Synopsis der europäischen Orthoptera [fourth part]. *Lotos* **3** : 168-176.
- FINOT, A. 1893. Séance du 25 janvier, 1893. *Annls Soc. ent. Fr.* **62** : xviii-xli.
- FISCHER DE WALDHEIM, G. 1846. *Orthoptera Imperii Rossici*. iii + 443 pp., 37 pls. Moscow.
- FOERSTER, A. 1868. Synopsis der Familien und Gattungen der Ichneumonen. *Verh. naturh. Ver. preuss. Rheinl.* **25** : 135-221.
- GERMAR, E. F. 1817. *Reise nach Dalmatien und in das Gebiet von Ragusa*. xii + 323 pp., 11 pls. Leipzig & Altenburg.
- GERSTAECKER, A. 1869. Beitrag zur Insekten-Fauna von Zanzibar. II. Orthoptera et Neuroptera. *Arch. Naturgesch.* **35** (1) : 201-223.

- GIGLIO-TOS, E. 1907. Spedizione al Ruwenzori di S. A. R. Luigi Amedeo di Savoia Duca degli Abruzzi. XVI. Ortotteri nuovi (diagnosi preventiva). *Boll. Musei Zool. Anat. comp. R. Univ. Torino* **22** (556) : 1-3.
- GRANT, H. J. 1964. A revision of the genera *Ceraia* and *Euceraia*, with notes on their relationship to *Scudderia* (Orthoptera; Tettigoniidae; Phaneropterinae). *Proc. Acad. nat. Sci. Philad.* **116** : 29-117, 228 figs.
- GRIFFINI, A. 1906. Ortotteri raccolti da Leonardo Fea nell'Africa occidentale. I. Hetrodidi, Conocephalidi, Meconemidi, Pseudophyllidi, Mecopodidi e Fanerotteridi. *Annali Mus. civ. Stor. nat. Genova* (3) **2** : 358-397.
- 1908a. Phasgonouridae Africane del R. Museo di Storia Naturale di Bruxelles. 5. Phaneropteridae 1. *Mém. Soc. ent. Belg.* **15** : 74-86.
- 1908b. Phasgonouridae Africane del R. Museo di Storia Naturale in Bruxelles. 6. Phaneropteridae 2. *Mém. Soc. ent. Belg.* **15** : 201-226.
- 1909. Note sopra alcune Phasgonouridae del Congo. *Annals Soc. ent. Belg.* **53** : 9-28, 1 fig.
- HARZ, K. 1957. *Die Gerafflügler Mitteleuropas.* xxiii + 494 pp., 254 figs., 20 pls. Jena.
- 1960. Gerafflügler oder Orthoptera (Blattodea, Mantodea, Saltatoria, Dermaptera). *Tierwelt Dtl.* **46**, xii + 232 pp., 566 figs.
- IMMS, A. D. 1957. *A general textbook of entomology.* 9th edn, revised by O. W. Richards and R. G. Davies. x + 886 pp., 606 figs. London.
- JACOBY, M. 1893. Descriptions of some new species of Eumolpidae and Halticidae from Africa (Gaboon). *Entomologist* **26** (Suppl.) : 97-102.
- JANNONE, G. 1936. Nuovi contributi alla conoscenza della fauna delle isole italiane dell'Egeo. Pt. 5. Studio bio-ecologico e sistematico dell'Ortottero-fauna con notizie sui Blattoidei, Mantoidei e Fasmoidei. *Boll. Lab. Zool. gen. agr. R. Scuola Agric. Portici* **29** : 47-248, 48 figs., 1 pl.
- KARABAĞ, T. 1964. Some new species and new records of Tettigoniidae (Orthoptera) from Turkey. *Communs Fac. Sci. Univ. Ankara* (C) **13** : 36-55, 40 figs.
- KARNY, H. H. 1910. Zoologische und anthropologische Ergebnisse einer Forschungsreise im westlichen und zentralen Südafrika ausgeführt in den Jahren 1903-1905 mit Unterstützung der Kgl. Preussischen Akademie der Wissenschaften zu Berlin von Dr. Leonhard Schultze. 4. Systematik und Tiergeographie. Orthoptera (s. str.). *Denkschr. med.-naturw. Ges. Jena* **16** : 35-90, 1 pl.
- 1915. Ergebnisse der Forschungsreise des Herrn Dr. Adalbert Klaptoz nach Französisch Guinea. Orthoptera und Oothecaria. *Zool. Jb. (Syst. Geogr. Biol.)* **40** : 119-146.
- 1926. On Malaysian katydids (Tettigoniidae). Represented in the collections of the F.M.S. Museum (Kuala Lumpur) and the Raffles Museum (Singapore). *J. fed. Malay St. Mus.* **13** : 69-156, 48 figs., 2 pls.
- 1927. Revision der Gryllacriden des Zoologischen Institutes in Halle a. S., sowie einiger Tettigoniiden-Typen von Burmeister und Giebel. *Z. Naturw.* **88** : 1-14.
- KARSCH, F. 1887a. Verzeichniss der von Herrn Waldemar Belck 1885 im Damaralande gesammelten Orthopteren. *Ent. Nachr.* **13** : 39-46.
- 1887b. Zwei neue ostafrikanische Phaneropteriden. *Ent. Nachr.* **13** : 52-54.
- 1888. *Weissenbornia*, eine neue Orthopteren- (Phaneropteriden-) Gattung aus Deutsch-Westafrika. *Ent. Nachr.* **14** : 65-67.
- 1889. Orthopterologische Beiträge III. *Berl. ent. Z.* **32** : 415-464, 1 pl.
- 1890a. Orthopterologische Mittheilungen 4. Ueber Phaneropteriden. *Ent. Nachr.* **16** : 57-62.
- 1890b. Neue westafrikanische, durch Herrn Premierlieutenant Morgen von Kribi eingesendete Orthopteren. *Ent. Nachr.* **16** : 257-276, 4 figs.
- 1890c. Verzeichniss der von Herrn Dr. Paul Preuss auf der Barombi-Station in Deutsch-Westafrika 1890 gesammelten Locustodeen aus den Familien der Phaneropteriden, Mekonemiden und Gryllakriden. *Ent. Nachr.* **16** : 353-369, 4 figs.

- KARSCH, F. 1892a. Uebersicht der von Herrn Dr. Paul Preuss auf der Barombi-Station in Kamerun gesammelten Locustodeen. *Berl. ent. Z.* **36** : 317-346, 7 figs.
 —— 1892b. Verzeichniss der von Herrn Dr. Paul Preuss im Kamerungebirge erbeuteten Orthopteren. *Berl. ent. Z.* **37** : 65-78, 3 figs.
 —— 1893. Die Insecten der Berglandschaft Adeli im Hinterlande von Togo (Westafrika) nach dem von den Herren Hauptmann Eugen Kling (1888 und 1889) und Dr. Richard Büttner (1890 und 1891) gesammelten Materiale. I. Apterygota, Odonata, Orthoptera Saltatoria, Lepidoptera Rhopalocera. *Berl. ent. Z.* **38** : 1-266, 35 figs., 6 pls. 1 map.
 —— 1896. Neue Orthopteren aus dem tropischen Afrika. *Stettin. ent. Ztg* **57** : 242-359, 44 figs.
 KIRBY, W. F. 1900. Notes on the collection of African Phasgonuridae formed by Mr. W. L. Distant in the Transvaal etc., with descriptions of two new species. *Ann. Mag. nat. Hist.* (7) **6** : 211-217.
 —— 1906. *A synoptic catalogue of Orthoptera* **2**, Pt. 1, viii + 562 pp. London.
 —— 1909. Ruwenzori Expedition reports. 8. Orthoptera. *Trans. zool. Soc. Lond.* **19** : 63-66.
 KRAUSS, H. 1890. Beitrag zur Kenntniß westafrikanischer Orthopteren. 2. Orthopteren der Guinea-Inseln São Thomé und Rolas, gesammelt von Prof. Dr. Richard Greeff. *Zool. Jb. (Syst. Geogr. Biol.)* **5** : 647-668, 1 pl.
 —— 1893. Vorläufige Diagnosen der neuen Arten und Varietäten von Oran. *Jh. Ver. vaterl. Naturk. Württ.* **48** : xcvi.
 —— 1901. Beitrag zur Kenntniß der Orthopteren Deutsch-Südwestafrikas. *Verh. zool.-bot. Ges. Wien* **51** : 281-293.
 LA BAUME, W. 1911. Beitrag zur Kenntniß der aethiopischen Orthopteren. *Stettin. ent. Ztg* **72** : 308-326, 19 figs.
 LE PELETIER DE SAINT-FARGEAU, A. L. M., & SERVILLE, J. G. A. 1828. *Encyclopédie méthodique. Histoire naturelle. Entomologie, ou histoire naturelle des crustacés, des arachnides et des insectes* **10**, 833 pp. Paris.
 MATSUMURA, S., & SHIRAKI, T. 1908. Locustiden Japans. *J. Coll. Agric. imp. Univ. Tokyo* **3** : 1-80, 2 pls.
 MORALES AGACINO, E. 1947. Notas sobre ortopteroideos de Ifni y Sáhara Español. *Eos, Madr.* **23** : 241-283, 4 figs.
 —— 1950. Apuntes sobre los Phasmidae y Tettigoniidae marroquíes del Instituto Español de Entomología. *Eos, Madr.* **26** : 157-196, 17 figs.
 MÜLLER, J., & HENLE, J. 1837. Sitzung der physicalisch-mathematischen Klasse (31. Juli). *Ber. Verh. Akad. Wiss. Berl.* **1837** : III-118.
 RAGGE, D. R. 1956a. A revision of the genera *Phaneroptera* Serville and *Nephoptera* Uvarov (Orthoptera : Tettigoniidae), with conclusions of zoogeographical and evolutionary interest. *Proc. zool. Soc. Lond.* **127** : 205-283, 131 figs.
 —— 1956b. A revision of the genus *Oxyconus* Chopard, 1935 (Orthoptera : Tettigoniidae). *Proc. R. ent. Soc. Lond. (B)* **25** : 183-190, 23 figs.
 —— 1957. A revision of the genus *Tropidonotacris* Chopard, 1954 (Orthoptera : Tettigoniidae). *Proc. R. ent. Soc. Lond. (B)* **26** : 119-122, 7 figs.
 —— 1960a. The Acrometopae of the Ethiopian Region: a revision, with notes on the sexual dimorphism shown by the group (Orthoptera : Tettigoniidae). *Bull. Br. Mus. nat. Hist. (Ent.)* **8** : 269-333, 148 figs.
 —— 1960b. Further notes on the genus *Phaneroptera* Serville, with a revised key to the species (Orthoptera : Tettigoniidae). *Proc. zool. Soc. Lond.* **134** : 237-250, 56 figs.
 —— 1961a. A revision of the genus *Ducetia* Stål (Orthoptera : Tettigoniidae). *Bull. Br. Mus. nat. Hist. (Ent.)* **10** : 171-208, 91 figs.
 —— 1961b. Further notes on the genus *Lamecosoma* Ragge, with a description of a new species (Orth. Tettigoniidae). *Eos, Madr.* **37** : 215-219, 5 figs.
 —— 1962a. A revision of the genera *Drepanophyllum* Karsch and *Stenamblyphyllum* Karsch (Orth. Tettigoniidae). *Eos, Madr.* **38** : 299-309, 21 figs.

- RAGGE, D. R. 1962b. A revision of the genera *Phlaurocentrum* Karsch, *Buettneria* Karsch and *Leiodontocercus* Chopard (Orthoptera : Tettigoniidae). *Bull. Br. Mus. nat. Hist. (Ent.)* **13** : 3-17, 32 figs.
- 1964. A revision of the genus *Tylopsis* Fieber (Orthoptera : Tettigoniidae). *Bull. Br. Mus. nat. Hist. (Ent.)* **15** : 297-322, 52 figs.
- RAMBUR, M. P. 1838-1842. *Faune entomologique de l'Andalousie* **2**, 336 pp., 18 pls. Paris.
- RAMME, W. 1951. Zur Systematik, Faunistik und Biologie der Orthopteren von Südost-Europa und Vorderasien. *Mitt. zool. Mus. Berl.* **27** : 3-431, 134 figs., 39 pls., 3 maps.
- REHN, J. A. G. 1914. *Wissenschaftliche Ergebnisse der Deutschen Zentral-Afrika-Expedition 1907-1908 unter Führung Adolf Friedrichs, Herzogs zu Mecklenburg. Band V. Lief. 1. Orthoptera I. Mantidae, Phasmidae, Acrididae, Tettigoniidae und Gryllidae aus dem Zentral-Afrikanischenseengebiet, Uganda und dem Ituri-Becken des Kongo.* Pp. 1-223. Leipzig.
- SAUSSURE, H. DE, & PICTET, A. 1897-1899. Fam. Locustidae. *Biologia cent.-am.* (Zool.) Insecta. Orthoptera **1** : 285-458, pls. 14-22.
- SCHAUM, H. 1853. Gesammsitzung der Akademie (15 December). *Ber. Verh. Akad. Wiss. Berl.* **1853** : 747-780.
- SCHULTHESS SCHINDLER, A. DE. 1898. Orthoptères du pays des Somalis, recueillis par L. Robecchi-Brichetti en 1891 et par le Prince E. Ruspoli en 1892-93. *Annali Mus. civ. Stor. nat. Genova (2)* **19** : 161-216, 1 pl.
- SERVILLE, J. G. A. 1831. Revue méthodique des insectes de l'ordre des Orthoptères [second part]. *Annls Sci. nat.* **22** : 134-166.
- 1838*. *Histoire naturelle des insectes. Orthoptères.* xviii + 776 pp., 14 pls. Paris.
- SJÖSTEDT, Y. 1901. Beiträge zur Kenntniss der Insektenfauna von Kamerun. 8. Locustodeen aus Kamerun und Kongo. *Bih. K. svenska VetenskAkad. Handl.* **27** (4, 3) : 1-45, 4 pls.
- 1909. *Wissenschaftliche Ergebnisse der schwedischen zoologischen Expedition nach dem Kilimandjaro, dem Meru und den umgebenden Massaisteppen deutsch-Ostafrikas 1905-1906 unter Leitung von Prof. Dr. Yngve Sjöstedt.* 17. Orthoptera. 6. Locustodea. Pp. 125-148, 1 pl. Stockholm.
- 1912. Zur Orthopterenfauna des Kamerungebirges. *Ark. Zool.* **7** (37) : 1-30, 4 figs., 3 pls.
- 1913. Neue Orthopteren aus Ost- und Westafrika nebst einigen anderen zugehörigen Formen. *Ark. Zool.* **8** (6) : 1-26, 3 pls.
- 1929. Voyage au Congo de S.A.R. le Prince Léopold de Belgique 1925. Orthoptera. 2. Phasgonurodea. *Revue Zool. Bot. afr.* **17** : 38-43, 3 figs.
- SPINOLA, M. 1839. Essai sur les Fulgorelles, sous-tribu de la tribu des Cicadaires, ordre des Rhynogotes [first part]. *Annls Soc. ent. Fr.* **8** : 133-337.
- STÅL, C. 1856. Orthoptera cursoria och Locustina från Cafferlandet. *Öfvers. K. VetenskAkad Förh.* **13** : 165-170.
- 1873. Orthoptera nova descripsit. *Öfvers. K. VetenskAkad. Förh.* **30** (4) : 39-53.
- 1874. Recensio Orthopterorum. *Revue critique des Orthoptères décrits par Linné, De Geer et Thunberg* **2**, 121 pp. Stockholm.
- 1876. Bidrag till södra Afrikas Orthopter-fauna. *Öfvers. K. VetenskAkad. Förh.* **33** (3) : 29-76.
- STEINHEIL, E. 1873. Symbolae ad historiam Coleopterorum Argentiniae meridionalis, ossia enumerazione dei coleotteri raccolti dal prof. P. Strobel nell'Argentina meridionale, e descrizione delle specie nuove. *Centuria II. Atti Soc. Ital. Sci. nat.* **15** : 554-578.
- STEINMANN, H. 1966. New *Phaneroptera* Serv. and *Nephoptera* Uv. species (Orthoptera: Tettigoniidae). *Acta zool. hung.* **12** : 409-417, 29 figs.
- STRAND, E. 1921. Neue Namen dreier Insekten-gattungen. *Ent. Z. Frankf. a. M.* **34** : 106.

* This work is dated "1839" and this is often cited as the date of publication. The work was, however, published in December, 1838 (teste Sherborn, C. D., 1922, *Index Animalium* (2) **1** : xxxiii).

- THOMSON, J. 1857. Description de Cérambycides nouveaux ou peu connus de ma collection. *Archs ent. Paris* **1** : 291-320.
- UVAROV, B. P. 1924. Notes on the Orthoptera in the British Museum. 3. Some less known or new genera and species of the subfamilies Tettigoniinae and Decticinae. *Trans. ent. Soc. Lond.* **1923** : 492-537, 1 fig., 1 pl.
- 1934. Entomological expedition to Abyssinia, 1926-7: Orthoptera of the families Mantidae Gryllidae, Tettigoniidae, and Acrididae. *J. Linn. Soc. (Zool.)* **38** : 591-614, 10 figs.
- 1936. Studies in the Arabian Orthoptera. I. Descriptions of new genera, species and subspecies. *J. Linn. Soc. (Zool.)* **39** : 531-554, 5 figs., 2 pls.
- 1940a. Twenty-eight new generic names in Orthoptera. *Ann. Mag. nat. Hist.* (11) **5** : 173-176.
- 1940b. Twenty-four new generic names in Orthoptera. *Ann. Mag. nat. Hist.* (11) **6** : 112-117.
- 1952. Studies in the Arabian Orthoptera. III. New genera, species and subspecies collected by the Anti-Locust Missions. *J. Linn. Soc. (Zool.)* **42** : 176-194, 55 figs.
- UVAROV, B. P., & POPOV, G. 1957. The Saltatorial Orthoptera of Socotra. *J. Linn. Soc. (Zool.)* **43** : 359-389, 38 figs., 1 pl., 1 map.
- WALKER, F. 1869. Catalogue of the specimens of Dermaptera Saltatoria in the collection of the British Museum **2** : 225-423. London.
- 1870. Catalogue of the specimens of Dermaptera Saltatoria in the collection of the British Museum **3** : 425-604. London.
- WERNER, F. 1929. Wissenschaftliche Ergebnisse einer zoologischen Forschungsreise nach Westalgerien und Marokko. II. Dermapteren und Orthopteren aus Westalgerien und Marokko. *Sber. Akad. Wiss. Wien (I)* **138** : 167-188, 7 figs., 4 pls.
- 1932. Orthopteren aus Marokko und Westalgerien. *Zool. Anz.* **100** : 113-127, 2 figs.
- ZACHER, F. 1917. Die Geredflügler Deutschlands und ihre Verbreitung. viii + 287 pp., 1 map. Jena.
- ZEUNER, F. E. 1936a. The subfamilies of Tettigoniidae (Orthoptera). *Proc. R. ent. Soc. Lond. (B)* **5** : 103-109.
- 1936b. The Recent and fossil Tympanophorinae (Tettigoniidae, Saltatoria). *Trans. R. ent. Soc. Lond.* **85** : 287-302, 2 pls.
- 1939. Fossil Orthoptera Ensifera. xiii + 321 pp., 80 pls. London.

INDEX

Valid names are in roman type, invalid names in italics.

- | | |
|--------------------------------------|------------------------------------|
| acaciae, 92 | ampla, 97 |
| accrana, 80 | amplectens, 82 |
| acuticercus, 83 | <i>amplipennis</i> , 87 |
| adeps, 94 | amydra, 96 |
| adicra, 84 | <i>Anerota</i> , 92 |
| aequatorialis, 84 | angustipennis, Arantia, 80 |
| <i>affinis</i> , 87 | angustipennis, Debrona, 82 |
| afra, 81 | angustipennis, Leiodontocercus, 89 |
| africana, Phaneroptera, 92 | angustipennis, Mangomaloba, 89 |
| africana, Poreuomena, 94 | Anisotochra, 80 |
| africana, Symmetropleura, 77, 78, 95 | annulifemur, 94 |
| albida, 92 | antinorii, 89 |
| algerica, 90 | apertus, 91 |
| amabilis, 96 | apicalis, 81 |

INDEX

- arabica*, 85
Arantia, 80
asymmetrica, 91
Atlasacris, 80
atrolineata, 80
attenuata, 97
Azamia, 80

banzyvilleana, 96
belcki, 82
bicolor, 85
bilineolata, 97
biplagiata, 80
bipunctata, 88
biramosa, *Ducetia*, 83
biramosa, *Zeuneria*, 98
blandus, 90
Bongeia, 81
borrei, 90
breviala, 84
brevicauda, 81
brevicollis, 85
brevipes, 80
brevis, *Odontura*, 91
brevis, *Tylopsis*, 97
brunneri, *Conchotopoda*, 78, 82
brunneri, *Melidia*, 89
Bueacula, 81
Buettneria, 81
Büttneria, 81

calaharica, 95
camerata, 82
Cameronia, 77, 78, 95
Camptoxiphae, 79
canaliculata, 85
capensis, 91
capreola, 81
carinata, *Tropidonotacris*, 96
carinatus, *Plangiodes*, 94
Catoptropteryx, 81
centralis, 98
Centrophorus, 96
cereris, 85
cervina, 82
Cestromoecha, 78, 81
chelocerca, 83
clavigera, 92
cloetensi, 91
coi, 97
compressa, 85
concava, 87
Conchotopoda, 82
condylus, 89

confluens, 97
contradti, 89
conspersa, 93
continua, 97
cooperi, 92
cornigera, 81
corrosa, 83
corrosifolium, 83
Corycomima, 82
Corymeta, 82
costata, 84
cothurnata, 85
crassicauda, 82
crassipes, 78, 81
crosskeyi, *Ducetia*, 84
crosskeyi, *Horatosphaga*, 87
crypterius, 84
cuspidata, 85

Dannfeltia, 82
Dapanera, 82
darlingi, 85
Debrona, 82
denticulata, 83
dilutum, 95
dimidiata, 86
diminuta, *Eurycorypha*, 85
diminuta, *Horatosphaga*, 87
Diogena, 83
Dioncomena, 83
dispar, 97
Dithela, 83
doriae, 80
Drepanophyllum, 83
dubia, 97
Ducetia, 83
duponti, 94

eburneiguttata, 81
eidmanni, 82
elegans, 89
elgonis, 87
elongata, *Horatosphaga*, 87
elongata, *Phaneroptera*, 92
Enochletica, 84
Epiphlebus, 84
Euanerota, 92
Eulioptera, 84
Eupantolepta, 87
Eurycorypha, 84
Euthyphlebia, 86
excelsior, 80
extensipes, 81

- fallax, 85
 fasciata, 80
 fatidica, 80
 fausta, 83
 feana, 86
 fenestrata, 96
 fissa, 97
 flavescens, 82
 foraminata, 94
 forcipata, 94
 fragilis, 92
 furcatum, 83
 fuscopunctata, 84

Gabonella, 85
Gabonia, 85
 gabunensis, 80
 galbana, 86
Gelatopoea, 85
Gelatopoia, 85
 genuteres, 82
 gestri, 80
 gladiator, 94
Goetia, 86
Gonatoxia, 86
 gracilipes, 80
 gracilis, *Horatospaga*, 87
gracilis, *Locusta*, 97
 gracilis, *Phaneroptera*, 92
gracilis, *Plegmatoptera*, 87
 gracilis, *Tylopsis*, 97
 grallatoria, 82
 graminea, 93
 grandis, *Pronomapyga*, 95
 grandis, *Tropidonotacris*, 96
 graueri, 95
Gravenreuthia, 86
 guttatypes, 81

 haasi, 91
habetata, 82
 hamuligera, 90
Harpocnepha, 86
 herbacea, 94
 heteromorpha, 87
Himertia, 86
Himertula, 86
hoechneli, 87
Horatospaga, 86
 hydatinoptera, 80

 immaculata, 86
 immaculipennis, 81
 inclusa, 87

 inconspicua, 93
indica, 92
 inermis, 88
inhama, 97
 insularis, 93
 irregularis, *Dapanera*, 82
 irregularis, *Tylopsis*, 97
Itokiia, 88
Ivensia, 88

Japygophana, 88

kandti, 87
karschi, *Pachypyga*, 87
 karschi, *Plangia*, 94
Karschia, 83
 karschiana, *Harpocnepha*, 86
 karschiana, *Pardalota*, 91
kasindina, 88
Keniacola, 87
 kenyensis, 89
Kevaniella, 88
 kilimandjarica, *Monticolaria*, 90
kilimandjarica, *Pantolepta*, 87
Kuwayamaea, 83

Lamecosoma, 88
 laminata, 89
 laminifera, 94
 lamottei, *Poreuomena*, 94
lamottei, *Tylopsis*, 97
 latevittatum, 93
latifolia, 80
latifrons, 90
latipennis, *Catoptropteryx*, 81
latipennis, *Mangomaloba*, 89
leggei, 87
Leiodontocercus, 89
lenzi, 93
leptocerca, 82
leptocnemis, 80
leptomorpha, 84
Leptophyes, 89
lesnei, 91
Letana, 89
liliifolia, 97
linearis, 87
liouvillei, 91
lobatipes, 94
lobatum, 93
lobulata, 96
lobulipennis, 86
longicerca, *Eulioptera*, 84

- longicercus, *Zeuneria*, 98
longipennis, 97
 longipes, *Horatsphaga*, 87
 longipes, *Tetraconcha*, 96
 longispina, 92
 loosi, 84
 lurida, 92

 macrocerca, 84
 maculata, *Gonatoxia*, 86
 maculata, *Peropyrrhicia*, 92
 maculatum, *Phlaurocentrum*, 93
 maculiceps, 81
 maculipennis, 81
 maculosa, 92
 magna, *Horatsphaga*, 87
 magna, *Phaneroptera*, 92
 magnus, *Oxyecous*, 91
 malleus, 89
 mammisignum, 80
 manca, 80
Mangomaloba, 89
marginalis, 92
marginata, 97
margineguttata, 97
 marmorata, *Arantia*, 80
 marmoratum, *Drepanophyllum*, 83
 maroccana, 91
 massaiae, 92
 mecopodoides, 93
 media, *Eurycorypha arabica*, 85
 media, *Horatsphaga*, 87
 melanopeza, 98
 melanotus, 80
 melica, 90
Melidia, 89
 meruensis, *Horatsphaga*, 87
 meruensis, *Monticolaria*, 90
meruensis, *Tylopsis*, 97
Meruterrana, 89
 microptera, 91
Milititsa, 90
Miltinobates, 90
 minima, 92
 minor, 95
 modulata, 90
moghrebica, 91
 montana, 85
Monteiroa, 90
 monticola, 89
Monticolaria, 90
 montivaga, 87
Morgenia, 90
multivenosa, 88

 mundamensis, 81
mutica, 85
Mylocentrum, 90
Myrmecophana, 84

 nana, *Dannfeltia*, 82
 nana, *Phaneroptera*, 92
natalensis, 93
nebulosa, 94
neutralipennis, 81
nigropunctata, 93
nigrospinosa, 81
nomima, 87
nuda, 87

 obesa, 97
obscuripes, 97
Odontura, 90
ornata, 83
ornatipes, 85
orthocnemis, 80
ostentatrix, 84
Otiaphysa, 82
ovalipennis, 80
Oxyecous, 91

Pachypyga, 86
Pachypygiana, 87
Pantolepta, 86
parallela, 87
Paranerota, 92
Parapyrrhicia, 91
Pardalota, 91
 parva, *Conchotopoda*, 82
 parva, *Ducetia*, 84
 parva, *Phaneroptera*, 93
Paura, 83
 peculiaris, 80
peeli, 88
 peloti, 88
Peronura, 92
Peropyrrhicia, 92
perpulchra, 97
Phaneroptera, 92
Phaneroptila, 93
Phaula, 93
Phaulula, 93
Phlaurocentrum, 93
Physocorypha, 93
 plana, 95
Plangia, 93
Plangiodes, 94
Plangiola, 94

- Plangiopsis*, 94
plasoni, 91
Plegmatia, 87
Plegmatoptera, 86
Poecilogramma, 94
politurata, 93
Poreuomena, 94
praestantissima, 98
praeusta, 97
Preussia, 94
producta, 95
Pronomapyga, 95
proserpinae, 85
Prosphega, 95
Pseudisotima, 83
Pseudopyrrhizia, 95
pulchra, 91
punctata, *Ducetia*, 84
punctata, *Pseudopyrrhizia*, 95
puncticollis, 81
punctipennis, *Ducetia*, 84
punctipennis, *Eurycorypha*, 85
punctulata, *Catoptropteryx*, 81
punctulata, *Phaneroptera*, 93
punctulata, *Tylopsis*, 97
Pyrrhicia, 89
Pyrrhizia, 89

quadridentata, 91
quadripunctata, *Phaneroptera*, 92
quadripunctata, *Telaea*, 84

ramulosa, 84
rectifolia, 80
rectiloba, 83
reducta, 85
regina, 80
regularis, 87
reimeri, 91
reticulata, *Eulioptera*, 84
reticulata, *Plegmatoptera*, 87
reticulosa, 84
retinervis, 80
Rhegmatopoda, 78, 82
rivae, 88
robusta, 87
roseata, 92
royi, 89
rubrescens, 97
rubricornis, 90
ruspolii, *Epiphlebus*, 84
ruspolii, *Horatosphaga*, 88

sagitta, 84
samburu, 88
saturata, 86
scalaris, 96
schoutedeni, 94
Schubotzcris, 95
Schlthessinia, 87
scotti, 92
surra, 80
securifera, 85
semiconchata, 94
serrifera, 88
signatipennis, 81
silvarum, 88
simillima, 85
simplicinervis, 80
smaragdina, 96
somali, 88
somaliensis, 90
sparsa, 92
spathulifera, 90
spinosus, 97
spinulicauda, 91
spinulosa, *Arantia*, 80
spinulosa, *Eulioptera*, 84
spinulosa, *Eurycorypha*, 85
splendens, 95
Stenacropteryx, 81
Stenamblyphyllum, 95
stenoxypha, 91
stichyrata, 96
stigmosum, 90
strangulata, 85
striatifemur, 94
stuhlmanni, 88
stylata, 85
stylifera, 88
subcarinata, 92
sudanensis, 85
superba, *Dioncomena*, 83
superba, *Pardalota*, 91
Symmetropleura, 77, 78, 95

Tapeina, 95
Tapiena, 95
teknicus, 91
Telaea, 83
Tellidea, 96
tenuicerca, 93
tenuipes, 81
tenuis, 88
terniensis, 91
Terpnistria, 96
Tetraconcha, 96

- tetrasticta*, 92
Thaumatoxenia, 87
tigrina, 80
tihamae, 96
Trigonocorypha, 96
trochlearis, 88
Tropidonotacris, 96
Tropidophrys, 96
tuberculata, 96
tuberous, 93
turbatum, 93
Tylopsis, 96
- ugandana*, 80
uncinata, 88
undulatus, 91
unimaculata, 94
uvarevi, 91
- varia*, 85
velicauda, 85
- versicolor*, 91
vicaria, 97
vicina, 88
villiersi, *Plangia*, 94
villiersi, *Tylopsis*, 97
viridis, 88
vitriala, 84
vittata, 97
vosseleri, 88
Vossia, 97
- Weissenbornia*, 98
wilverthi, 94
- zanzibarica*, *Letana*, 89
zanzibarica, *Parapryrrhicia*, 91
zebrata, *Eurycorypha*, 85
zebrata, *Terpnistria*, 96
Zeuneria, 98

